USANA Technical Bulletin

Disclaimer: The information provided in this Technical Bulletin is strictly educational. It may not be used to promote USANA products, nor is it intended as medical advice. For diagnosis and treatment of medical disorders, consult your health care professional. When there are references to third party websites, addresses, and/or phone numbers, USANA, Inc. makes no claim, actual or implied, regarding the content or validity of the information obtained from such sources. This Technical Bulletin may be copied and freely distributed only if all text remains intact and unchanged.

Pantothenic Acid (Calcium Pantothenate)

Technical Background

- Pantothenic acid (sometimes referred to as vitamin B₅) is an essential water-soluble nutrient widely synthesized by plants and many bacteria, but required in the diets of vertebrate animals, including humans.¹
- The principal active form of pantothenic acid is coenzyme A (CoA).² Mitochondria may be the final site of CoA synthesis, since 95% of this coenzyme is found in mitochondria.¹
- As a component of coenzyme A, pantothenic acid is essential for the production of energy from carbohydrate, fat and protein. It plays a pivotal role in energy metabolism and the production of ATP, the cell's principal energy currency.^{2,3}
- Pantothenic acid and coenzyme A are involved in the synthesis of many other biochemicals, including amino acids, fatty acids, proteins, cholesterol, steroid hormones, vitamin A, vitamin D, hemoglobin, cytochrome molecules, glutathione,⁴ and vitamin B₁₂.⁵
- Coenzyme A also contributes an acetyl group to the important neurotransmitter acetylcholine.²
- Pantothenic acid may have a protective effect against oxidative damage and ultraviolet light exposure.^{6,7} Studies have also found that increasing intake of pantothenate improves wound healing.⁸
- Pantothenic acid is widely distributed in foods, where about 85% occurs as coenzyme A. Because of this broad availability, acute pantothenic acid deficiencies are rare in humans. Where such deficiencies have occurred (generally in conjunction with severe malnutrition), they have been associated with such symptoms as headache, fatigue, insomnia, numbing of the toes, and painful, burning sensations in the hands and feet.^{1,5}

Sources and Recommended Intake

- The name "pantothenic acid" is derived from the Greek word "pantos," meaning "widespread." This is a reference to the fact that the nutrient is found in most foods. The richest dietary sources include meats, fish, eggs, broccoli, avocados, and legumes.¹
- Pantothenic acid is unstable to heating, acids, and bases. As such, significant amounts can be destroyed during food preparation. Cooking is reported to destroy 15-50% of the pantothenic acid in meat, and studies have shown that 37-78% of the pantothenic acid in fruits and vegetables can be lost during processing.⁹
- The Recommended Dietary Allowance (RDA) has been set at 5mg per day for adults; most people consume 4-7 mg per day.¹⁰

- People with diabetes mellitus, inflammatory bowel diseases, or chronic alcoholism typically have increased requirements for pantothenic acid.⁵
- Calcium pantothenate and sodium pantothenate are the forms of pantothenic acid most often found in nutritional supplements.
- High oral doses of pantothenic acid or calcium pantothenate (up to 10 grams per day for several weeks) do not appear to be toxic to humans.¹¹

Abstract

Slyshenkov VS, Piwocka K, Sikora E, Wojtczak L. Pantothenic acid protects jurkat cells against ultraviolet lightinduced apoptosis. Free Radic Biol Med. 2001 Jun 1;30(11):1303-10. Human leukemic T lymphocytes (Jurkat cells) were induced to undergo apoptosis by brief irradiation with ultraviolet C light (254 nm). This was accompanied by accumulation of lipid peroxidation products in the form of conjugated dienes, a decrease of total glutathione content, and a shift of its redox state towards the oxidized form. Preincubation of the cells with 1 mM pantothenate resulted in a significant elevation of total glutathione content of the cells, reaching its maximum level, 160% of the control, after 3 h. Similar increase was observed after preincubation with 5 mM N-acetylcysteine, a known precursor of glutathione. Both pantothenic acid and N-acetylcysteine alleviated the ultraviolet-induced decrease of glutathione content, diminished lipid peroxidation, and partly protected the cells against apoptosis produced by ultraviolet irradiation.

References

¹ Plesofsky-Vig N. Pantothenic Acid. In Ziegler EE, Filer LJ (eds). Present knowledge in Nutrition. Washington DC: ILSI Press. 1996. p 236-244.

² Lehninger AL. 1975. Biochemistry. New York:Worth Publishers. 1975. 1104 pp.

³ Depeint F, Bruce RW, Shangari N, Mehta R, O'Brien PJ. Mitochondrial function and toxicity: Role of the B vitamin family on mitochondrial energy metabolism. 2006. Chemico-Biological Interactions. 163(1-2):94-112.

⁴ Slyshenkov VS, Dymkowska D, Wojtczak L. Pantothenic acid and pantothenol increase biosynthesis of glutathione by boosting cell energetics. 2004. FEBS Letters 569(1-3):169-172.

⁵ Groff JL, Gropper SS, Hunt SM. Advanced Nutrition and Human Metabolism. New York: West Publishing Company. 1995. 575 pp.

⁶ Wojtczak L, Slyshenkov VS. Protection by pantothenic acid against apoptosis and cell damage by oxygen free radicals--the role of glutathione. Biofactors. 2003;17(1-4):61-73.

⁷ Slyshenkov VS, Moiseenok AG, Wojtczak L. Noxious effects of oxygen reactive species on energy-coupling processes in Ehrlich ascites tumor mitochondria and the protection by pantothenic acid. Free Radic Biol Med. 1996;20(6):793-800.

⁸ Weimann BI, Hermann D. Studies on wound healing: effects of calcium D-pantothenate on the migration, proliferation and protein synthesis of human dermal fibroblasts in culture. Int J Vitam Nutr Res. 1999 Mar;69(2):113-9.

⁹ Tahiliani AG, Beinlich CJ. Pantothenic acid in health and disease. Vitam Horm 1991;46: 165-228.

¹⁰ Institute of Medicine. Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline (1998). National Academies Press: Washington, D.C.

¹¹ Bender DA. Nutritional Biochemistry of the Vitamins. New York: Cambridge University Press. 1992. p 341-59.