

THE USES AND BENEFITS OF
ALPHA LIPOIC ACID

by

REBECCA MAULDIN

The Uses and Benefits of Alpha Lipoic Acid

Alpha lipoic acid, also called thioctic acid, is a sulfur-containing fatty acid compound that acts as a coenzyme and an antioxidant. It is called a “universal antioxidant” because of its potent ability to neutralize a wide range of free radicals. Recent studies suggest that alpha lipoic acid may improve situations in the body involving free radicals, such as heart disease and cancer, and other disorders associated with inflammation and aging. The uses and benefits of alpha lipoic acid are numerous, making it an important nutrient for maintaining and regaining health in the 21st century.

Although many studies have been done on alpha lipoic acid, the research is still in its early stages. So far, the research is showing that this potent antioxidant may be one of the most important supplements people could take to improve their health. With the ever increasing amount of toxic chemicals and environmental pollutants in the world, the human body is now exposed to more sources of free radicals than ever before. Free radicals cause harmful chemical reactions that damage cells and attack tissue in the body causing premature aging, inflammation, allergies, and enzyme dysfunction. As a result, a person becomes more susceptible to chronic diseases such as diabetes, atherosclerosis, and cancer. As we look closer at the uses and benefits of alpha lipoic acid, I will first explain what it is and when it was discovered. Then we will look at its biological activities in the body and how it benefits the body through several mechanisms. Finally,

we will see how alpha lipoic acid has been proven to be useful for particular illnesses and diseases.

Alpha lipoic acid was first isolated in 1951 by biochemist Lester Reed Ph.D. Dr. Reed purified a very small amount of alpha lipoic acid crystals from a sample of liver weighing a total of 250 pounds.¹ Alpha lipoic acid is a naturally occurring compound that is produced in small amounts by plants, animals, and humans.² It was found to be a co-factor for several mitochondrial enzyme complexes that are involved in energy production, similar in function to many of the B vitamins.³ But unlike the B vitamins, alpha lipoic acid is not classified as a vitamin because it is able to be synthesized by the body. Alpha lipoic acid is found in abundance in animal tissues with high metabolic activity, such as kidney, heart, and liver, and to a lesser extent in fruits and vegetables. Non-animal sources are spinach, broccoli, tomatoes, peas, Brussels sprouts, and rice bran. All alpha lipoic acid supplied by the diet is transported in the bloodstream to tissues and incorporated into cells. It is then transferred into the mitochondria. Studies in mammals have shown the alpha lipoic acid supplied by the diet does not supply enough for purposes such as incorporation into enzyme complexes.⁴ Hence, these sources of alpha lipoic acid are thought to provide very little free alpha lipoic acid into circulation. It is only through supplementation that alpha lipoic acid reaches potentially therapeutic levels.

Alpha lipoic acid works on the cellular level to help produce energy in the body. To do this, it acts as a coenzyme in the energy cycle of the cell, called the citric acid cycle. The conversion of food to energy is driven by a series of chemical reactions that

are set in motion by enzymes. Alpha lipoic acid takes a part in this multienzyme process by preparing the fuel for the mitochondria, the power plants of the cell, and plays a vital role in mitochondrial electron transport reactions required for metabolizing glucose into adenosine triphosphate, or ATP, for cellular energy production.⁵

Alpha lipoic acid is a medium chain fatty acid with two sulfur atoms. The reduced form is known as dihydro-lipoic acid and the oxidized form is known as alpha lipoic acid. Alpha lipoic acid also contains an asymmetric carbon, which means that there are two possible optical isomers that are mirror images of each other: R-lipoic acid and S-lipoic acid. R-lipoic acid is the natural form of alpha lipoic acid, while S-lipoic acid is the synthetic form. It is the natural form of alpha lipoic acid that has been found to have the greatest benefit to the body. Researchers have found that synthetic alpha lipoic acid does not improve ATP synthesis in isolated cells like the natural form does. The natural R-form was shown to increase membrane fluidity and transport, while the S-form was much less effective in doing so.⁶ The reduced form has been found to exert a number of antioxidant and neuro-protective actions that are not seen in alpha lipoic acid. While both forms are able to scavenge a number of free radicals, only R-dihydro-lipoic acid has been shown effective against superoxide and peroxy reactive oxygen species.⁷

The oxidized and reduced forms of R-lipoic acid made up a redox couple. Oxidation reduction or “redox” reactions involve the transfer of an electron from a donor to an acceptor. This transfer changes the reduced form to the oxidized form when the donor loses an electron. When an acceptor gains an electron, it changes from its oxidized

form to its reduced form. It is through this cycle of reduction to R-dihydro-lipoic acid and oxidation back to R-lipoic acid that beneficial processes take place, such as the ability to provide cysteine, an amino acid that is critical for glutathione production.⁸

It is the R-form of alpha lipoic acid that occurs naturally in foods, covalently bound to lysine in proteins, such as beef liver. Because the content of alpha lipoic acid in foods is limited, dietary supplements are likely as much as 1000 times greater than the amounts that could be obtained through the diet. In the United States, alpha lipoic acid is available without a prescription and has been available since the 1990's.⁹ Most alpha lipoic acid supplements contain a 50/50 mixture of R-lipoic acid and S-lipoic acid. (There are a few companies that offer the R-form of alpha lipoic acid for sale, but it more expensive than alpha lipoic acid.) Direct comparisons of the bioavailability of the 50/50 mixture and R-lipoic acid supplements have not been published. However, a few studies have found that after oral dosing with the 50/50 mixture of the R-lipoic acid and the S-lipoic acid, peak plasma concentrations of the R-form were found to be 40-50% higher than the S-form, suggesting that R-lipoic acid is better absorbed than S-lipoic acid.^{10 11}

Because alpha lipoic acid provides benefits to so many basic cellular functions, it has a great future as a supplement for many diseases. It may be one of our best defenses against disease and premature aging. A closer look at the beneficial properties of alpha lipoic acid will demonstrate its vital role in the body.

Alpha lipoic acid has a wide range of benefits. The most important benefit is its role as an antioxidant. In 1988, the antioxidant properties of alpha lipoic acid were discovered. It was shown to have powerful antioxidant abilities, equal to that of coenzyme Q 10, vitamin C, and vitamin E.¹² Free radicals are highly unstable electrons that are a normal byproduct of oxidative metabolism. When they get out of control, these electrons can cause extensive damage to lipid membranes, organelles, and to DNA itself. Unlike other antioxidants, alpha lipoic acid has the unique ability to neutralize free radicals within aqueous and lipid regions of the cells, as well as in intracellular and extracellular environments.¹³ This ability allows alpha lipoic acid to be easily transported across cellular membranes to neutralize free radicals. Vitamin C is only able to protect the watery portions of cells from free-radical attack, and vitamin E is only able to protect the fatty membranes. Alpha lipoic acid, on the other hand, has the ability to neutralize free radicals that occur in both the watery and fatty regions of the cell. Alpha lipoic acid is capable of neutralizing a wide variety of free radicals, such as singlet oxygen, superoxides, peroxy and hydroxyl radicals, hypochlorite, and peroxynitrite.¹⁴ Other antioxidants, like vitamin C, vitamin E, or glutathione can only neutralize one kind of free radical. Researchers call alpha lipoic acid the “universal antioxidant” because it can neutralize such a wide variety of free radicals.

Not only is alpha lipoic acid able to neutralize free radicals, it is also able to recycle or regenerate several other important antioxidants, including vitamin C and glutathione. When an antioxidant scavenges a free radical, it becomes oxidized in the process and is not able to scavenge additional free radicals until it has been reduced. The

reduced form of alpha lipoic acid, dihydro-lipoic acid, is able to reduce the oxidized antioxidants, enabling them to be useful again.¹⁵ Increasing glutathione levels is very important because it is one of the body's most important intracellular antioxidants.

Glutathione is a compound that is composed of 3 amino acids: cysteine, glutamic acid, and glycine. It protects the body from free radical damage and plays a leading role in the detoxification and elimination of potential carcinogens and toxins in the body.

Glutathione conjugation is one of the primary mechanisms for eliminating xenobiotics in the liver.¹⁶ Animal studies have shown that glutathione synthesis and cellular glutathione levels are significantly lower in aged animals than in younger animals, leading to a decreased ability of aged animals to respond to toxin exposure and oxidative stress.¹⁷

Alpha lipoic acid, when fed to the aged animals, has been found to increase glutathione synthesis in cultured cells and in animal tissues.¹⁸ In a study with rats, oral dosing of 150 milligrams per kilogram of body weight per day for eight weeks significantly increased the glutathione levels in the blood and in the liver.¹⁹ Increases in glutathione levels that alpha lipoic acid provides are not only from the reduction of oxidized glutathione, but also from the synthesis of glutathione.²⁰ The availability of cysteine inside a cell determines its rate of glutathione synthesis. Alpha lipoic acid, by its reduction to dihydro-lipoic acid and oxidation back to alpha lipoic acid, has the ability to continuously provide cysteine, an important amino acid for glutathione production. It reduces extracellular cystine to cysteine and increases the uptake of cysteine into the cell, increasing glutathione production.²¹

Alpha lipoic acid has been found to reduce the formation of glycosylated end products, or AGEs.²² AGEs are formed when proteins react with sugars, and this process increases the risk of cardiovascular disease by oxidizing LDL cholesterol and making blood vessels tough and inflexible. Alpha lipoic acid can also affect the left ventricle of the heart, reducing its ability to pump blood into the circulatory system, thereby increasing blood pressure. Glycosylated proteins are also unable to bind to receptors on liver cells to signal the cessation of cholesterol manufacturing. This causes the body to continue to produce too much cholesterol. Alpha lipoic acid stops these processes from happening by inhibiting glycation at the starting point.²³ In a study with rats, the effect of alpha lipoic acid was investigated on the formation of AGEs. The rats were given a glucose solution to drink for 4 weeks, combined with a normal chow diet or a diet supplemented with alpha lipoic acid. The chronic administration of the glucose resulted in a 29% increase in blood pressure, 30% increase in glycemia, 286% increase in insulinemia, 408% increase in insulin resistance index, and a 63% increase in AGE content in the aorta. The rats that were given alpha lipoic acid did not have the increases in blood pressure, insulin levels, insulin resistance, and aorta AGE content. By preventing AGE formation, alpha lipoic acid prevented the development of hypertension and hyperglycemia in the rats.²⁴

Another benefit of alpha lipoic acid is its ability to chelate, or bind, with heavy metals. Both forms of alpha lipoic acid have been shown to form complexes with manganese, zinc, cadmium, lead, cobalt, nickel, and iron ions.²⁵ In one study, an intraperitoneal injection of 25 milligrams of alpha lipoic acid given to rats for seven days

was able to significantly alter the oxidative stress caused by lead toxicity.²⁶ In another study, alpha lipoic acid was demonstrated to be able to protect rat hepatocytes from cadmium toxicity by preventing decreases in total glutathione and increases in lipid peroxidation.²⁷ These protective effects have also been seen in rats that had depleted levels of glutathione prior to cadmium exposure.²⁸ A study with rats examined the effects of alpha lipoic acid on arsenic toxicity. Arsenic toxicity caused decreases in glutathione peroxidase in the brain of the rats, decreased activity of superoxide dismutase, and decreased catalase activity. All regions of the brain also showed increased levels of reactive oxygen species and other intracellular oxidants and an increase in lipid peroxidation levels. Rats that were given alpha lipoic acid simultaneously with the arsenic did not demonstrate the deficits in antioxidant enzyme activities or the increase in lipid peroxidation. Alpha lipoic acid was able to prevent arsenic-induced changes in the antioxidant defense system of the rats.²⁹ Alpha lipoic acid has also been found to protect against chromate-induced oxidative stress in mouse livers.³⁰ While other chelators such as dimercaptosuccinic acid (DMSA) and 2,3-dimercapto-1-propanesulfonic acid (DMPS) are unable to cross the blood-brain barrier and remove heavy metals from the brain, alpha lipoic acid is particularly suited to this purpose as it can penetrate both the blood-brain barrier and the cell membrane.^{31 32}

Protection from irradiation is another benefit of alpha lipoic acid. Studies have evaluated the effects of alpha lipoic acid on radiation injury in both animals and humans. In mice, it has been demonstrated that alpha lipoic acid, but not dihydro-lipoic acid, was able to protect against radiation injury to cells involved in blood cell formation.

Administration of alpha lipoic acid increased the survival rate of irradiated mice from 35% in untreated animals to 90% in treated animals.³³ In a Russian study involving Chernobyl radiation victims, alpha lipoic acid was shown to greatly reduce free radical damage caused by the radiation exposure. A group of 16 children given 400 milligrams of alpha lipoic acid for 28 days had significant reduction in both white blood cell free radical activity and urinary excretion of radioactive isotopes compared to a control group. Alpha lipoic acid lowered levels of chemiluminescence, a marker for lipid peroxidation, to levels of children who were not exposed to radiation. Combining vitamin E with alpha lipoic acid further reduced chemiluminescence to below normal levels. Vitamin E alone did not have this effect. Kidney and liver function was also normalized by the alpha lipoic acid treatment.³⁴

Alpha lipoic acid has demonstrated the ability to protect the liver from poisons and the ability to regenerate the liver if it has already been damaged. Some of the first human clinical studies using alpha lipoic acid were carried out by Fredrick C. Bartter and Burton M. Berkson from the National Institutes of Health in the 1970's. Dr. Berkson has since used it successfully for the treatment of many chronic liver diseases, such as viral hepatitis and autoimmune hepatitis. Alpha lipoic acid's ability to regenerate the liver is demonstrated in a study of three people with liver damage from Hepatitis C. Through the use of alpha lipoic acid, selenium, and silymarin, all three were able to avoid liver transplants and regained liver function.³⁵ Alpha lipoic acid infusions were used in the treatment of potentially deadly amanita mushroom poisoning in 75 patients between 1974

and 1978. Eighty nine percent of the patients recovered after receiving the lipoic acid infusion, while only 10-50 percent of the patients recovered without intervention.³⁶

Alpha lipoic acid has a beneficial effect on cholesterol and lipid levels. One study, which explored the effect of antioxidant capacity on blood lipid metabolism and lipoprotein lipase (LPL) activity of rats fed a high-fat diet, found that alpha lipoic acid improved the antioxidant capacity and activity of LPL and reduced blood lipids significantly.³⁷ In a study with mice fed a high fat diet, supplementation with alpha lipoic acid caused a decrease in lipid peroxidation, plasma cholesterol, triglycerides, and low-density lipoprotein cholesterol, and an increase in high-density lipoprotein cholesterol.³⁸ The effect of alpha lipoic acid on lipid peroxidation and lipid levels has been studied in the blood, liver, and kidneys of young and aged rats. Lipid levels were considerably higher in the aged rats compared with the young rats. Alpha lipoic acid was administered intraperitoneally for 7 and 14 days. It was shown to prevent the elevated lipid levels seen in the aged rats.³⁹

Alpha lipoic acid has been found to prevent calcium oxalate crystals or stones from forming in and damaging the kidneys. Hyperoxaluria is the excretion of an excessive amount of oxalate in the urine and is the major risk factor for the formation of urinary calcium oxalate stones. In a study with rats, alpha lipoic acid and eicosapentaenoic acid (EPA) were shown to reduce tubular damage and to decrease the markers of crystal deposition substantially in the animals. The EPA and alpha lipoic

acid was effective at inhibiting stone formation and was able to protect the kidneys from damage caused by oxalate toxicity.⁴⁰

Nuclear factor kappa beta (NF-kappa B) is a transcription factor. Transcription factors are messengers found inside cells, and they carry information from the cytoplasm to the nucleus. They activate or inhibit the production of certain proteins or enzymes, which then carry out a particular cell function. Nuclear factor kappa beta is an inflammatory messenger which turns on inflammation. By stabilizing nuclear factor kappa beta transcription factors in the body, alpha lipoic acid can regulate a number of genes that are related to inflammation and cell cycle control, which are involved in the pathology of diabetes, atherosclerosis, and cancer.⁴¹ Th1- and Th2-mediated immune system cells identify and respond to oxidative stress with certain cell membrane receptors. These receptors initiate a cascade of events that eventually activate the master transcription factor NF-kappa B. NF-kappa B is then able to bind to DNA and affect the rate of transcription of certain deleterious genes. Because of this, NF-kappa B plays a significant role in the regulation of inflammatory-induced gene function. High doses of alpha lipoic acid have been shown to inhibit the activation of NF-kappa B when added to cell culture.⁴² Because of this ability, alpha lipoic acid has been used for the treatment of various cancers for which no effective treatments exist. Dr. Berkson used intravenous alpha lipoic acid, low dose naltrexone, and other oral antioxidants to achieve long term survival of a patient with metastasized pancreatic cancer.⁴³ Another case study, published in 2007, described the complete reversal of B-cell lymphoma in a patient using

less than one month of intravenous alpha lipoic acid and six months of low dose naltrexone.⁴⁴

Alpha lipoic acid also displays antihyperglycemic effects.⁴⁵ It has been shown to aid in increasing glucose uptake in skeletal muscles as well as improving insulin-stimulated glucose disposal.⁴⁶ Alpha lipoic acid works against insulin resistance by increasing the permeability of cell membranes, which is decreased in hyperglycemia.⁴⁷ A study was done on the effect of alpha lipoic acid on insulin sensitivity in people with type 2 diabetes mellitus. Twelve patients were given an oral 600 milligram tablet of alpha lipoic acid twice a day, one tablet in the morning and the other in the evening, 30 minutes before a meal, for a period of 4 weeks. Insulin sensitivity was measured using a manual hyperinsulinaemic euglycaemic clamp technique. The 4-week treatment with alpha lipoic acid significantly improved glucose utilization in the type 2 diabetic patients. In half of the patients, the amount of glucose metabolized after the treatment reached a level considered normal for healthy people.⁴⁸

Another significant benefit of alpha lipoic acid lies with the mitochondria. Mitochondria are cellular organelles that oxidize dietary proteins, fats, and carbohydrates into a useable form of energy, called adenosine triphosphate, or ATP. Mitochondria provide energy for basic metabolic processes, and their decay with age impairs cellular metabolism and leads to decline of cellular function. Mitochondrial dysfunction appears to contribute to some of the loss of function accompanying aging because aged tissue uses oxygen inefficiently, which impairs ATP synthesis, resulting in increased free

radical production.⁴⁹ The high amount of oxidants not only damages the mitochondria, but may eventually cause inadequate energy production and the general metabolic decline evident in aging. In a study using old rats, research found that supplementing with 0.5% R-lipoic acid for 2 weeks partially reversed the age-associated loss of mitochondrial function, the increase in oxidative stress, and the damage and decline in general metabolic activity. The supplemented rats were significantly more active, which further shows that R-lipoic acid acts physiologically to increase general metabolic activity.⁵⁰

Systemic inflammation plays an important role in several diseases. Alpha lipoic acid has been found to have significant benefits on inflammation due to its ability to down regulate pro-inflammatory cytokines. When free radicals disrupt the normal structure of proteins and the genetic information encoded in DNA, this damage always results in the release of potent chemical messengers called cytokines. Cytokines trigger the inflammatory response in tissues, signaling immune and inflammatory cells to swarm the affected area. Alpha lipoic acid inhibits prostaglandin E2 production by inhibiting COX 2 activity.⁵¹ Another study investigated the prophylactic effect of alpha lipoic acid on systemic inflammation in mice. The mice were divided into 3 groups: control, systemic inflammation, and alpha lipoic acid treated mice with systemic inflammation. Mice with systemic inflammation had higher levels of reactive oxygen species in lymphocytes, hepatocytes, and astrocytes, than in the control mice. However, the mice receiving the alpha lipoic acid did not show elevated levels of the reactive oxygen species. Systemic inflammation also increased levels of lipid peroxidation and decreased levels of superoxide dismutase, catalase, glutathione peroxidase, and glutathione. These

levels were reduced to almost normal levels when the mice were treated with alpha lipoic acid.⁵²

Stress is a stimulus that can alter physiological homeostasis and can cause many physiological changes in the body. The ability to cope with stress is a crucial determinant of health and disease.⁵³ There have been studies that have indicated that long-term exposure to stress has a detrimental effect on several brain functions and increases the production of free radicals in the body.⁵⁴ Stress has been shown to cause depletion of the glutathione-based antioxidant defense system of the body and to decrease the level of vitamin C in the body.⁵⁵ In a study with rats, the effect of stress on lipid peroxidation and antioxidant enzymes in the retina and in the brain were specifically examined. The researchers found that stress caused a significant decrease in brain copper/zinc superoxide dismutase and in the levels of glutathione peroxidase in the brain and retina. Rats given alpha lipoic acid while being exposed to stress were protected from lipid peroxidation by the antioxidant properties of alpha lipoic acid.⁵⁶ Alpha lipoic acid might provide indirect benefit when cortisol levels are high because it can partially restore the hydrocortisone-induced suppression of helper T-cell activity.⁵⁷ It has also been shown to prevent the accumulation of catecholamines in cardiac tissue in response to stress and to enhance the elimination of catecholamine degradation products.⁵⁸

Due to the broad range of health benefits alpha lipoic acid is able to provide, it has been used in the treatment of many chronic degenerative diseases. The biological activities of alpha lipoic acid make it an important supplement with the potential for both

preventing and treating disease. It has been shown to have a beneficial effect on many different diseases, from cataracts to cancer.

Because cataracts are associated with reduced antioxidant activity in the lens of the eye, alpha lipoic acid has been shown in animal studies to inhibit cataract formation. Research has shown that lenses with cataracts have dramatic decreases in reduced glutathione, as much as 81 percent, when compared with normal lenses.⁵⁹ Results of one rat study showed that 25 milligrams per kilogram of body weight of alpha lipoic acid protected 60 percent of the animals from cataract formation. The major biochemical changes in the lens associated with the protective effect of alpha-lipoic acid were increases in glutathione, ascorbate, and vitamin E levels.⁶⁰ The researchers in this study found that the reduced form of alpha lipoic acid, R-dihydro-lipoic acid, reached higher concentrations in the rat lenses than the synthetic form of alpha lipoic acid.⁶¹ This data demonstrated that R-lipoic acid was more effective in maintaining glutathione levels and protecting the lens from damage than the S-lipoic acid.

Low levels of glutathione may also contribute to the pathological processes in the development of glaucoma. Supplementation of alpha lipoic acid has been shown to increase glutathione in red blood cells and in the lacrimal fluid of patients with glaucoma.⁶² In a Russian trial of alpha lipoic acid, 45 patients with stage I and II glaucoma were given either 75 milligrams or 150 milligrams for 2 months. A control group of glaucoma patients was given only local hypotensive medication. Improvement was seen in 45 percent of those patients receiving alpha lipoic acid.⁶³

The antioxidant properties of alpha lipoic acid make it beneficial for reversing the aging process. There is a declining ability with age to respond to oxidative stress. Oxidative stress is a condition that arises when our cells cannot adequately destroy the free radicals that are a normal byproduct of cellular processes. Oxidative stress also results from exposure to toxic chemicals such as pesticides, hydrocarbons, smog, alcohol, and prescription drugs. During the aging process, there is an increase of mitochondrial decay,⁶⁴ oxidant production, and oxidative damage to important molecules.⁶⁵ These processes may in part be responsible for aging as well as age-associated degenerative diseases such as cancer and atherosclerosis.⁶⁶ In one study, researchers from the University of California at Berkeley took liver cells from aging rats and measured their energy level, free radical level, and the ability of the cells to recycle vitamin C. The aged rats were 3 times less active than the young rats. The free radicals were 5 times higher in the aged rats, the energy production was very low, and the ability to recycle vitamin C in the aged rats was only half of the levels of the young rats. After 2 weeks on R- lipoic acid, the free radicals in the aged rats decreased, their vitamin C levels rose, and their energy levels rose significantly. Glutathione levels also increased when the aged rats were given alpha lipoic acid. Improvement in metabolic activity as well as lowering oxidative stress and the damage caused by aging were significantly improved by the use of alpha lipoic acid.⁶⁷ With age, glutathione levels naturally decline, making older animals more vulnerable to both free radicals and other environmental toxins. Another study with rats examined the effect of alpha lipoic acid and acetyl-L-carnitine on age-related decline in metabolic function and oxidative stress. The combination was able to

partially reverse the decline in average mitochondrial membrane potential and significantly increased the ambulatory activity in both the young and old rats. Levels of vitamin C in liver cells, which were very low in the aged rats, were restored to the level seen in young rats when they were given alpha lipoic acid and acetyl-L-carnitine.⁶⁸

Photoaging of the facial skin has shown improvement with topical applications of alpha lipoic acid. In one study on 33 women with a mean age of 54.4 years, half of the face was treated twice a day for 12 weeks with a cream containing 5% alpha lipoic acid and the other half with a control cream. The methods used to assess the improvement were self-evaluation by the women, clinical evaluation, photographic evaluation, and laser profilometry. All four methods of assessment showed a significant improvement on the side of the face treated with the cream containing alpha lipoic acid compared to the placebo-treated half of the face.⁶⁹

Alpha lipoic acid has been used to improve mental function and might be a successful therapy for Alzheimer's disease and other related dementias. Mitochondria are known to lose efficiency with age due to the oxidation of proteins, lipids, DNA, and RNA.⁷⁰ Age-related decay of mitochondrial function can be partially reversed in older animals by treatment with dihydro-lipoic acid. In one study, older rats receiving the reduced form of alpha lipoic acid showed signs of reduced lipid peroxidation and improved memory. Electron microscope studies in the hippocampus region of the brain showed that the alpha lipoic acid reversed age-associated mitochondrial structural decay.⁷¹ In a study with aged beagle dogs, a daily supplement of alpha lipoic acid and

acetyl-L-carnitine for a period of 2 months significantly improved their learning ability on two landmark discrimination tasks compared to controls.⁷²

Alpha lipoic acid has been shown to have a variety of properties which can interfere with pathogenic principles of Alzheimer's disease. For example, alpha lipoic acid increases acetylcholine (ACh) production by activating choline acetyltransferase and increases glucose uptake, thus supplying more acetyl-CoA for the production of ACh. Alpha lipoic acid chelates redox-active transition metals, thus inhibiting the formation of hydroxyl radicals and also scavenges reactive oxygen species, thereby increasing the levels of reduced glutathione. Via the same mechanisms, a decrease of inflammatory processes is also achieved. Alpha lipoic acid can scavenge lipid peroxidation products such as hydroxynonenal and acrolein. The reduced form of alpha lipoic acid, dihydro-lipoic acid, is the active compound responsible for most of these beneficial effects.⁷³ In a study by German researchers, alpha lipoic acid was administered to 9 Alzheimer's patients for a year. The supplement led to a stabilization of cognitive functions as determined by constant scores in two neuropsychological tests.⁷⁴ And in a second study, researchers have shown that alpha lipoic acid protects cortical neurons from the toxic effects of two oxidative substances implicated in Alzheimer's disease.⁷⁵

Parkinson's disease, a neurodegenerative disorder, may also be improved by alpha lipoic acid. In an animal model, the pathogenic mechanisms that lead to neurodegeneration were studied by administering a neurotoxin to mice. This toxin was shown to activate apoptosis signal regulating kinase (ASK1) and the phosphorylation of

downstream kinases MKK4 and JNK. The administration of alpha lipoic acid with the neurotoxin abolished these effects and was able to afford neuroprotection.⁷⁶

Other chronic diseases that alpha lipoic acid has shown to be of benefit for are some of the most common chronic disease conditions: metabolic syndrome, diabetes, and cardiovascular disease. Metabolic syndrome is characterized by a group of risk factors that include obesity, high LDL cholesterol and triglycerides, high blood pressure, and insulin resistance. These risk factors increase the risk of developing cardiovascular disease and diabetes. In animal experiments, alpha lipoic acid has been shown to cause weight loss, ameliorate insulin resistance and atherogenic dyslipidemia, as well as lower blood pressure.⁷⁷ Alpha lipoic acid has been shown to have anti-obesity effects. There is a fuel sensor in the cell called AMP-activated protein kinase that is activated when cellular energy is depleted. Administration of alpha lipoic acid to rodents has caused a decrease in AMP-activated protein kinase activity, causing profound weight loss by reducing food intake and enhancing energy expenditure.⁷⁸ In a recent randomized, double-blind clinical trial, endothelium-dependent vasodilation of the brachial artery was increased by 67%, 44%, and 75% in groups receiving irbesartan (an angiotensin-blocking drug), lipoic acid, and irbesartan plus lipoic acid, respectively, compared to placebo. Treatment with irbesartan and lipoic acid was also associated with significant reductions in plasma levels of pro-inflammatory mediators, such as interleukin-6.⁷⁹

The most widely noted clinical application of alpha lipoic acid is in the treatment of diabetes and its complications. Acting as an antioxidant, R-dihydro-lipoic acid has

been shown to protect rat pancreatic islet cells from destruction by reactive oxygen species.⁸⁰ Lester Packer, an antioxidant authority, notes that alpha lipoic acid works against insulin resistance by increasing the permeability of cell membranes, which is decreased and prevents the uptake of glucose.⁸¹ High doses of alpha lipoic acid can improve glucose utilization in individuals with type 2 diabetes by increasing glucose uptake in skeletal muscles, as well as by enhancing insulin-stimulated glucose disposal.^{82 83} The increased burning of glucose may occur because alpha lipoic acid enhances glucose utilization by the mitochondria in the cells. In a small clinical trial, 13 patients with type 2 diabetes mellitus were given a single intravenous infusion of 1000 milligrams of alpha lipoic acid. The alpha lipoic acid improved insulin sensitivity by 50% compared to a placebo infusion.⁸⁴ Another study of 72 patients with type 2 diabetes found that oral administration of alpha lipoic acid at doses of 600 milligrams per day, 1200 milligrams per day, or 1800 milligrams per day improved insulin sensitivity by 25% after 4 weeks of treatment.⁸⁵ They found that there were not significant differences among the three doses of alpha lipoic acid, which indicated that 600 milligrams per day may be the maximum effective dose.⁸⁶ In a study of lean and obese patients with type 2 diabetes, treatment with 600 milligrams of alpha lipoic acid was found to prevent hyperglycemia-induced increases of serum lactate and pyruvate levels and was found to increase glucose effectiveness.⁸⁷ R-alpha lipoic acid may be more effective in improving insulin sensitivity than S-alpha lipoic acid, according to studies with animals. One study found that R-alpha lipoic acid increased glucose uptake by skeletal muscle of obese rats by 65%, while S-alpha lipoic acid only increased glucose uptake by 29%.⁸⁸

Alpha lipoic acid has been used extensively in Germany for the treatment of diabetic neuropathy, a type of nerve damage that causes pain, loss of sensation and weakness.⁸⁹ Both oral and intravenous alpha lipoic acid are approved for the treatment of diabetic neuropathy in Germany.⁹⁰ Three large-scale, double blind, placebo-controlled trials, called the Alpha-Lipoic Acid in Diabetic Neuropathy (ALADIN) studies, have looked at the neurological effects of alpha lipoic acid. The first ALADIN study found that alpha lipoic acid used for 3 weeks intravenously at 600 or 1,200 milligrams per day was superior to a placebo for reducing the symptoms of neuropathy. The 600 milligram dose had slightly better results with fewer side effects.⁹¹ The second ALADIN study examined nerve conduction parameters as well as Neuropathic Disability Score of 65 patients with type 1 or 2 diabetes. The trial lasted for 2 years, and patients were divided into three groups: the first group received 600 milligrams of alpha lipoic acid twice a day, the second group received 600 milligrams of alpha lipoic once a day, and the third group received a placebo. The study found that alpha lipoic acid resulted in significant improvement in some of the nerve conduction parameters, but not in the Neuropathic Disability Score compared to the placebo group.⁹² In the third ALADIN trial, 509 type 2 diabetics with peripheral neuropathy received either 600 milligrams of alpha lipoic acid intravenously for three weeks, followed by 600 milligrams orally three times daily for six months, or 600 milligrams of intravenous alpha lipoic acid for three weeks followed by a placebo for six months, or a double placebo. In the groups treated with alpha lipoic acid, there was a significant improvement in nerve function.⁹³ In another trial, known as the SYDNEY trial, 120 type 2 diabetics were given 600 milligrams of alpha lipoic acid intravenously or a placebo for five days a week for a total of 14 treatments. The group

receiving the alpha lipoic acid reported significant improvement in overall symptoms compared to the placebo.⁹⁴ A meta-analysis of four placebo-controlled trials that have been done (ALADIN I and III, SYDNEY and an unpublished trial) found that the same treatment with 600 milligrams per day of intravenous alpha lipoic acid for 3 weeks in all trials significantly reduced the symptoms of diabetic neuropathy, and the analysis also found continuous daily improvement in symptom scores beginning on the eighth day of treatment.⁹⁵

The oral form of alpha lipoic acid in the treatment of neuropathy is also beneficial, although studies show it may not be quite as useful as the intravenous form. In one trial, 24 type 2 diabetic patients were randomly assigned 600 milligrams of alpha lipoic acid three times a day or a placebo for three weeks. Significant improvement in symptoms of pain, burning, and numbness were reported in the group receiving the alpha lipoic acid.⁹⁶ In one of the longest controlled trials using oral alpha lipoic acid therapy, 299 diabetic patients with peripheral neuropathy were given either 1200 milligrams per day of alpha lipoic acid, 600 milligrams per day of alpha lipoic acid, or a placebo. The length of the study was 2 years and by that time only 65 of the original participants were included in the final analysis. Of those, the patients taking either amount of alpha lipoic acid showed significant improvement in electrophysiological tests of nerve conduction compared with those who took the placebo.⁹⁷

Cardiac autonomic neuropathy is another neurological complication of diabetes. It occurs in as many as 25% of diabetic patients.⁹⁸ In a trial of 72 patients with type 2

diabetes, 800 milligrams per day of alpha lipoic acid resulted in a significant improvement in 2 out of 4 measures of heart rate variability when compared to a placebo.⁹⁹ Studies with mice have revealed that alpha lipoic acid may act as a protective agent for reducing cardiovascular complications of diabetes. Mice that were made diabetic and normal mice that served as controls were given a high fat/low cholesterol diet with or without alpha lipoic acid. At 20 weeks, the markers of oxidative stress were significantly lower in the mice receiving the alpha lipoic acid compared to those without it. The increase in cholesterol, atherosclerotic lesions, and the general deterioration of health caused by diabetes was completely prevented by alpha lipoic acid. There was also a reduction of blood glucose levels and the regeneration of insulin producing cells in the pancreas. The cells in the pancreas were also protected from damage.¹⁰⁰ Diabetics are also at high risk for vascular disease because endothelial function is often impaired in diabetic patients.¹⁰¹ In a randomized controlled trial, 58 patients diagnosed with metabolic syndrome were given 300 milligrams per day of alpha lipoic acid orally for 4 weeks. Using ultrasound to measure flow-mediated vasodilation, those receiving alpha lipoic acid improved 44% compared to the placebo group.¹⁰²

Delayed or impaired wound-healing is another complication of diabetes caused by the underlying factor of peripheral vascular dysfunction. Injuries often fail to heal because of high concentrations of pro-inflammatory cytokines present in the wound. Alpha lipoic acid has been shown to improve wound healing in studies with rats. In one study, diabetic rats were given alpha lipoic acid for 8 weeks. They were then subjected to abrasion wound formation. The wounds in the diabetic rats treated with alpha lipoic acid

healed more rapidly than the wounds in the untreated diabetic rats.¹⁰³ These findings suggest that alpha lipoic acid might help to improve the healing of wounds in diabetics. In a study with humans undergoing hyperbaric oxygen therapy, alpha lipoic acid supplementation was able to accelerate wound repair in patients affected by chronic wounds. Alpha lipoic acid, in combination with hyperbaric oxygen therapy, was able to down-regulate inflammatory cytokines and growth factors, promoting the healing process.¹⁰⁴

Alpha lipoic acid has been found to be useful for multiple sclerosis. Reactive oxygen species, or free radicals, play an important role in many of the events underlying multiple sclerosis. Reactive oxygen species are known to effect the migration of monocytes and cause dysfunction of the blood brain barrier. To infiltrate the central nervous system, monocytes must cross the blood brain barrier. They are then able to begin the process of demyelination and axonal damage. In a rat model for multiple sclerosis, acute experimental allergic encephalomyelitis, alpha lipoic acid prevented the development of the clinical signs of multiple sclerosis. Alpha lipoic acid inhibited the migratory capacity of the monocytes, and also stabilized the integrity of the blood brain barrier.¹⁰⁵ In one experiment with mice, feeding high doses of alpha lipoic acid was found to slow the progression of autoimmune encephalomyelitis, a model of multiple sclerosis.¹⁰⁶ The alpha lipoic acid inhibited the activity of an enzyme known as matrix metalloproteinase (MMP)-9 in these animals. When a small pilot study evaluated the safety of alpha lipoic acid in 30 people with relapsing of progressive multiple sclerosis,

treatment with 1200-2400 milligrams per day of oral alpha lipoic acid showed that it was well-tolerated and that there were decreases in serum MMP-9 levels.¹⁰⁷

Migraine headaches may be improved by treatment with alpha lipoic acid. In a study in Belgium, a randomized double-blind placebo-controlled trial of alpha lipoic acid was undertaken to see if alpha lipoic acid may be beneficial in migraine headache treatment. Five Belgian centers recruited 44 patients with migraine headaches. They received either a placebo or 600 milligrams of alpha lipoic acid per day for 3 months. Monthly migraine attacks were reduced during the treatment period in the alpha lipoic acid group compared to the placebo group. Within-group analyses showed a significant reduction in the frequency of migraine attacks, migraine days, and migraine severity in the patients treated with alpha lipoic acid for 3 months. No adverse effects were reported and this study indicates that alpha lipoic acid may be a beneficial migraine prophylaxis.¹⁰⁸

Alpha lipoic acid may be of benefit in cardiovascular disease. The risk factors of atherosclerosis, high cholesterol, smoking, diabetes mellitus, high homocysteine levels and hypertension, have one thing in common: the generation of oxidative stress.¹⁰⁹ Oxidative influences on LDL cholesterol cause an increase in atherogenicity by altering cell receptor uptake of these particles.¹¹⁰ The oxidized LDL is taken up by scavenger receptors on monocytes, smooth muscle cells and macrophages in a process leading to the accumulation of lipids and the formation of foam cells, an early feature of atherosclerotic plaques. Inflammatory events then occur within the newly formed lesion that further

generate peroxides, superperoxides and hydroxyl radicals within the endothelium, causing a cycle that damages the vasculature. Because of its ability to recycle antioxidants, alpha lipoic acid may protect against free radical LDL cholesterol oxidation and decrease the risk of cardiovascular disease.¹¹¹ In a study that was done with quail, alpha lipoic acid was shown to have a protective effect on the buildup of plaque in the arteries, a characteristic of atherosclerosis. The group receiving the alpha lipoic acid had 75% fewer atherogenic lesions than the control group. The researchers suggested that the alpha lipoic acid may guard against atherosclerosis by preventing the oxidation of LDL cholesterol and/or by recycling vitamin E.¹¹² Another study found that dietary alpha lipoic acid of 1 milligram per kilogram of food reduced the levels of total cholesterol and lipoproteins in the serum and aortic tissue of rabbits.¹¹³ In a more recent study, researchers found that supplementing mice with alpha lipoic acid reduced arterial lesion formation, triglycerides, blood vessel inflammation and weight gain, all of which are factors involved in the development of cardiovascular disease. They used apolipoprotein E-deficient and apolipoprotein E/low-density lipoprotein receptor-deficient mice, which are established models of human heart disease. The mice were divided into groups that received diets containing either a normal amount of fat or extra fat, with or without supplemental lipoic acid, for ten weeks. The amount of alpha lipoic acid was the human equivalent to 2 grams per day. The mice receiving the supplement had a significant reduction of atherosclerotic lesion formation in both groups eating varying amounts of fat. Supplemented mice also had 40 percent less weight gain and lower triglycerides in serum compared to those that did not receive the supplement. Less inflammation was also experienced in the supplemented mice.¹¹⁴ Researchers from Beijing University found

that alpha lipoic acid, because of its free-radical scavenging effects, is able to protect the myocardium from free-radical damage and decrease the incidence of malignant arrhythmias.¹¹⁵

Hypertension may be improved with alpha lipoic acid. In hypertensive patients, lower concentrations of antioxidants and superoxide dismutase activity have been documented.¹¹⁶ In one study, rats were given a diet supplemented with alpha lipoic acid for 9 weeks. After 9 weeks, results showed lowered systolic blood pressure, cytosolic $[Ca^{2+}]_i$, blood glucose and insulin levels, and tissue aldehyde conjugates.¹¹⁷ In another study, rats were given 10% D-glucose in their drinking water combined with either a normal chow diet or with an alpha lipoic supplemented diet, and they were compared with control rats during 3 weeks. The rats fed chronic glucose developed non-insulin dependant diabetes as reflected by an increase in both blood glucose and insulin levels. They also had a progressive increase in systolic blood pressure, an increased level of superoxide anion, and a lower plasma glutathione peroxidase activity level. However, supplementation of alpha lipoic acid in the diet of glucose fed rats prevented the rise in blood pressure, the increase in superoxide production, the decrease in plasma glutathione peroxidase, and the development of insulin resistance. This study suggests that the antihypertensive and hypoglycemic effects of alpha lipoic acid are associated with oxidative stress as reflected by the decrease in the basal superoxide anion production and by the preservation of the activity of glutathione peroxidase in the plasma of chronically glucose-treated rats.¹¹⁸

AMP-activated protein kinase (AMPK) acts as a fuel sensor in the cell and is activated when the energy in the cell is depleted. AMPK is important in the regulation of food intake and energy expenditure in the body. Alpha lipoic acid has been shown to decrease hypothalamic AMPK activity and cause profound weight loss in rodents by reducing food intake and improving energy metabolism. Through this mechanism, alpha lipoic acid exerts anti-obesity effects.¹¹⁹ Alpha lipoic acid is also able to enhance fatty acid oxidation and reduce lipid accumulation due to activating AMPK in skeletal muscle. In a study with rats, alpha lipoic acid was able to lower triglyceride accumulation in the skeletal muscle of obese rats and increase insulin-stimulated glucose disposal.¹²⁰ Obesity has been found to be associated with increased cardiovascular mortality.¹²¹ Endothelium-dependant vascular relaxation is impaired in obese rats and this impairment has been found to be associated with increased lipid accumulation and decreased AMPK activities in endothelial cells.¹²² This has also been seen to be an early event in the development of atherosclerosis.¹²³ In a study using obese rats, alpha lipoic acid was found to reduce food intake, normalize body weight, and normalize metabolic parameters, such as triglycerides and insulin. Alpha lipoic acid was found to improve vascular dysfunction and normalize AMPK activities in the endothelial cells.¹²⁴

Alpha lipoic acid may also be beneficial during an acute stroke. It is able to inhibit platelet and leukocyte activation and adhesion, reduce free-radical generation, and increase cerebral blood flow. Ischemia is the condition suffered by tissues and organs when deprived of blood flow. It is due mostly to the effects of inadequate nutrients and oxygen. Reperfusion injury refers to the tissue damage inflicted when blood flow is

restored after an ischemic period of more than about ten minutes. Ischemic-reperfusion injury in humans occurs in conditions such as stroke, cardiac arrest, subarachnoid hemorrhage or head trauma. Tissue damage is observed during reperfusion, which is primarily attributed to oxidative injury resulting from the production of free radicals. Several animal studies have shown that dihydro-lipoic acid, not alpha lipoic acid, has demonstrated a protective effect against ischemia reperfusion injury in the brain.^{125 126} However, in a gerbil animal study, it was the synthetic form of alpha lipoic acid that was found to protect against ischemia reperfusion injury. When animals were treated with alpha lipoic acid for seven days prior to the ischemia reperfusion injury, they had less change in locomotor activity and less damage to brain cells than the control animals.¹²⁷ In another study with rats, pretreatment with alpha lipoic acid before brain ischemia reduced mortality from 78% to 26% in the 24 hour period following the reperfusion. This pretreatment with alpha lipoic acid protected the brain from the loss of glutathione normally caused by ischemia reperfusion injury and significantly reduced the elevation in lipid peroxidation.¹²⁸ In another study in which hearts from rats fed alpha lipoic acid were analyzed, the alpha lipoic acid was found to preserve vitamin E in the heart tissue, to decrease lipid peroxidation, and to improve the recovery of the function of the left ventricle.¹²⁹ When alpha lipoic acid was studied on strokes induced in mice, administering a subcutaneous injection 1.5 hours before the stroke was induced was shown to reduce the infarct volume in middle cerebral artery occlusion. Treatment with alpha lipoic acid produced a significant reduction in lesion size at 24 hours and a significant improvement in neurological function at 24 hours. This study shows that taking alpha lipoic acid before the onset of a stroke can prevent some of the neurological

damage caused by the stroke. In this study, the mechanism of action of alpha lipoic acid appears to be due to its ability to substitute for glutathione. Free radical injury causes a depletion of cellular glutathione, leading to oxidation of protein thiols to disulfides and the loss of enzymes having thiol groups.¹³⁰

Alpha lipoic acid has been shown in animal models to have a beneficial effect on heavy metal poisoning. Heavy metals, particularly cadmium, arsenic, and mercury, cause cellular damage by harming the mitochondria, by inhibiting mitochondrial enzymes, by suppressing protein synthesis, and by producing free radicals.¹³¹ In one study, dogs and mice were completely protected from arsenite poisoning and mercury poisoning with high doses of alpha lipoic acid, but it was ineffective against lead and gold poisoning.¹³² Alpha lipoic acid can form a complex with arsenic that renders that arsenic nontoxic.¹³³ In studies using mice, alpha lipoic acid has been shown to prevent intestinal uptake of arsenic and to reduce the toxic effect of arsenic on enzyme inhibition.¹³⁴ Alpha lipoic acid has been shown to protect rat hepatocytes from cadmium toxicity at concentrations of 5 mM. It decreased cadmium uptake by the hepatocytes and normalized glutathione levels, increasing survival and cell viability despite cadmium toxicity.¹³⁵ In another study using rats, treatment with alpha lipoic acid completely prevented cadmium-induced lipid peroxidation of the heart, brain, and testes. Glutathione concentrations in the brain were 63% decreased in the rats treated with cadmium, but only 4% increased in rats treated with both cadmium and alpha lipoic acid.¹³⁶ Alpha lipoic acid can protect cells from lipid peroxidation caused by excess copper. It has been shown to bind to copper in human lipoproteins and to inhibit peroxidation of the lipoproteins.¹³⁷ Wilson's disease has been

successfully treated with alpha lipoic acid due to its ability to increase renal copper excretion and to normalize liver function.¹³⁸

High levels of iron, particularly in the brain, along with an increased level of unsaturated fatty acids, lead to increased levels of tissue peroxidation. The free radicals that are formed by reactions with iron in the central nervous system have been found to be suppressed by alpha lipoic acid.¹³⁹ The heavy metal platinum can cause renal toxicity. In experiments with rats with cisplatin-induced renal damage, large doses of alpha lipoic acid were able to restore the antioxidant enzyme activity levels to normal, increase glutathione levels, and decrease renal platinum content.¹⁴⁰ Lead toxicity has also been shown to be improved in animal models. In one study, an intraperitoneal injection of 25 milligrams of alpha lipoic acid was given to rats for seven days. It increased glutathione levels 207% and decreased malondialdehyde levels in the brain, kidneys, and red blood cells.¹⁴¹

Another use for alpha lipoic acid is for burning mouth syndrome and idiopathic dysgeusia, an altered perception of taste. In one study, 22 patients with idiopathic dysgeusia were given alpha lipoic acid for 2 months. The results showed significant symptom improvement compared to the placebo group.¹⁴² Another study looked at the effect of alpha lipoic acid on burning mouth syndrome. Forty-two patients with burning mouth syndrome were divided into 2 groups. One group was given 600 milligrams of alpha lipoic acid for 20 days, followed by 200 milligrams for 10 days. The control group was given a placebo for 30 days. Results showed significant improvement in two thirds

of the people receiving alpha lipoic acid, while only 15% of those taking the placebo had improvement. When those who had taken the placebo were switched to alpha lipoic acid, up to two thirds showed significant improvement as well.¹⁴³

As people age, they tend to produce excess levels of inflammatory cytokines such as interleukin-6, interleukin-1b, and tumor necrosis factor alpha (TNF-alpha). Cytokines are immune system-regulating chemicals. Pro-inflammatory cytokines can induce a state of chronic inflammation and can damage the bone matrix of the body. Studies have shown that alpha lipoic acid can inhibit the damage to human bone marrow cells caused by TNF-alpha. In one study, TNF-alpha caused an increase in intracellular reactive oxygen species and a reduction in cellular glutathione levels, leading to cell death. When the cells were pretreated with alpha lipoic acid, these changes induced by TNF-alpha were prevented. Therefore, alpha lipoic acid may have a therapeutic role in stopping bone loss associated with oxidative stress.¹⁴⁴ Another study found alpha lipoic acid to inhibit osteoclast differentiation by reducing NFK-kappa B DNA binding and to prevent bone loss.¹⁴⁵

Alpha lipoic acid may also play an important role in improving hearing loss. In a study with aging rats, alpha lipoic acid was found to improve cochlear function by improving mitochondrial function.¹⁴⁶ Damage to the ear is a side effect of many drugs and alpha lipoic acid may provide protection from these side effects. A study examined the use of alpha lipoic acid for hearing loss associated with cisplatin, a widely used anticancer drug. Cisplatin's clinical use is limited by the onset of severe side effects, one

of them being toxicity to the ear and deafness. Evidence suggests that free radicals are involved in the ototoxic effects of cisplatin and that the use of cisplatin results in depletion of glutathione and antioxidant enzymes (superoxide dismutase, catalase, glutathione-peroxidase and glutathione-reductase) in cochlear tissues, with a corresponding increase of the malondialdehyde levels.¹⁴⁷ When rats were given alpha lipoic acid along with cisplatin, it was shown to protect the antioxidant defense system of the ear.¹⁴⁸ Studies have also shown that alpha lipoic acid lessens nerve damage induced by aminoglycoside antibiotics. In experiments carried out with guinea pigs, alpha lipoic acid was found to reduce cochlear damage that the drug amikacin caused in the control group.¹⁴⁹

Exposure to high energy impulse noise caused by explosions can result in damage to the auditory system. Rats were used for a study which examined whether a short period of pre-exposure supplementation with antioxidants would protect the ear from the effects of noise. The rats given alpha lipoic acid for three days before being exposed to noise showed a much higher measurement of blood oxygenation than the controls. The alpha lipoic acid was able to protect the ear from free radical stress, antioxidant depletion, and hemoglobin oxidation that occurs after the exposure to high energy impulse noise.¹⁵⁰

Adriamycin, an anthracycline antibiotic, which is widely used as an anticancer drug in the treatment of various solid tumors, has been shown to induce reproductive abnormalities in males. In a study with rats, the effect of alpha lipoic acid was investigated on adriamycin-induced testicular toxicity. Adriamycin caused a significant

increase in DNA damage in the sperm, a significant decrease in the activities of steroidogenic enzymes, and decreased serum testosterone levels. However, treatment with alpha lipoic acid one day prior to adriamycin administration was able to maintain near normal steroidogenesis and spermatogenesis, thereby proving it to be an effective cell protectant.¹⁵¹ Adriamycin also causes a significant decrease in the activities of mitochondrial antioxidant enzymes, such as superoxide dismutase, glutathione peroxidase, and glutathione reductase, in the testis. But pre-treatment with alpha lipoic acid before exposure to adriamycin resulted in the ability to maintain near normal activities of the enzymes in the mitochondria in an experiment with rats.¹⁵² The oxidative stress caused by adriamycin also causes damage to the red blood cells, resulting in lowered hemoglobin and hematocrit levels and increased lipid peroxidation of the cell membrane. But when rats were given alpha lipoic acid before they were given the adriamycin, it was effective in counteracting the biochemical disturbances and toxic effects of the adriamycin.¹⁵³

Alpha lipoic acid is able to exert a protective effect on the liver when it is exposed to poisons and toxins. One study examined the protective effects alpha lipoic acid has on liver and kidney toxicity in rats that were given an oral toxic dose of acetaminophen. The rats were pretreated with 100 milligrams per kilogram of alpha lipoic acid and then given an oral toxic dose of acetaminophen. None of the animals pretreated with alpha lipoic acid died, and the elevation of nitric oxide production, reduction of glutathione peroxidase, and depletion of intracellular reduced glutathione levels in the liver and kidneys, were suppressed by the alpha lipoic acid. This treatment completely prevented

acetaminophen-induced mortality and inhibited oxidative stress in the liver and kidneys.¹⁵⁴ Another study looked at the action of alpha lipoic acid on the antioxidant system in rats with toxic hepatitis. Injections of alpha lipoic acid caused the levels of glutathione reductase and peroxidase to return to normal levels. Alpha lipoic acid was shown to regulate the manifestations of oxidative stress and the state of the glutathione antioxidant system.¹⁵⁵ A German study found alpha lipoic acid beneficial in the treatment of chronic liver diseases. When 89 patients with liver diseases (42 with chronic hepatitis, 25 with hepatic cirrhosis, 14 with hepatitis, and 8 with adiposis hepatica) were given oral and intravenous alpha lipoic acid, good results were achieved.¹⁵⁶

Researchers examined the protective effects of alpha lipoic acid on the oxidative damage caused by acrolein, a major toxicant in cigarette smoke, in human retinal pigment epithelial cells. Smoking is a cause of severe oxidative stress and has been shown to deplete ascorbic acid levels and cause oxidation of lipids and proteins.^{157 158} Three large studies have found a link between smoking and age-associated macular degeneration.¹⁵⁹ Of the six main toxic chemicals in cigarette smoke, acrolein has one of the highest hazard indexes and is much more toxic than formaldehyde and acetaldehyde, and can reach 80 μM in the respiratory tract fluid in smokers.¹⁶⁰ Acute exposure to acrolein in the retinal cells caused decreases in cell viability, mitochondrial potential, glutathione levels, antioxidant capacity, Nrf2 expression, and enzyme activity. Cells lost 79% and 87% of viability after ARPE-19 cells were exposed to 50 and 100 μM acrolein for 24 hours, whereas lower concentrations (1–25 μM) of acrolein had no apparent effect. When the cells were pretreated with alpha lipoic acid, they were protected from the oxidative

damage caused by acrolein exposure. Alpha lipoic acid was able to maintain the intracellular antioxidant levels, glutathione levels, and protect against mitochondrial damage. It has been estimated that the acrolein level that is in cigarette smoke is approximately 25 to 140 µg per cigarette.¹⁶¹ Because smoking is a habitual activity and the concentration of acrolein during smoking is less than that shown in the acute exposure seen in this study, the researchers also exposed the retinal cells to low doses of acrolein for 8 days or 32 days. They found that long term exposure caused similar toxicity in the cells.¹⁶²

Alpha lipoic acid has a remarkable effect on the liver. American physician, Dr. Burton Berkson, was one of the first doctors to conduct large human clinical studies on alpha lipoic acid. In the 1970's, he worked with Dr. Fredrick C. Bartter and other associates from the National Institutes of Health to investigate the use of alpha lipoic acid on liver damage. Intravenous alpha lipoic acid was administered to 79 people with acute and severe liver damage at several medical centers in the United States and 75 people recovered full liver function.¹⁶³ Dr. Berkson has since treated hundreds of patients with alpha lipoic acid for liver damage, autoimmune disease, cancer, and other diseases. His research and studies using alpha lipoic acid for hepatitis C have shown that it may be a low cost and efficacious alternative treatment program in combination with 2 other antioxidants. In Dr. Berkson's study, 3 patients with cirrhosis, portal hypertension and esophageal varices, secondary to a chronic hepatitis C infection, were given alpha lipoic acid, silymarin, and selenium. All 3 patients recovered quickly and their laboratory values improved markedly. They were able to avoid liver transplants and were able to

resume normal lives. A closer look at his program with 3 patients will show that alpha lipoic acid is highly beneficial to the health of the liver.

In modern medicine, there is no reliable therapy for chronic hepatitis C infection since interferon and antivirals work no more than 30% of the time. Liver transplantation is another option, but is not always successful due to the residual viruses that can infect the new liver. It is also extremely costly and can be a very painful and debilitating procedure. Dr. Berkson used the triple antioxidant combination of alpha lipoic acid, silymarin (milk thistle seed extract), and selenium in the treatment of 3 patients with chronic hepatitis C. These substances were chosen because of their ability to protect the liver from free radical damage, to increase the levels of other antioxidants in the body, and to interfere with viral proliferation. Each patient was given 600 milligrams of alpha lipoic acid, 900 milligrams of silymarin, and 400 micrograms of selenium per day. The patients were also prescribed B vitamins, vitamin C, vitamin E, and a mineral supplement. Dietary modifications that were suggested to the patients were to eat a diet that included at least 6 servings of vegetables and fruits, only 4 ounces or less of meat per meal, and 8 glasses of water per day. The patients were also instructed to reduce their stress levels and to take part in an exercise program.

The first patient, Mrs. M.P., was a 57-year-old woman who contracted hepatitis C after a blood transfusion during surgery 10 years ago. She did not have a healthy lifestyle at the time and did not eat a nutritious diet. She became very fatigued and nauseous about 5 years ago and was diagnosed with non-A, non-B hepatitis. After

treatment with conventional therapies, she continued to degenerate into a poorer state of health. She was diagnosed with chronic hepatitis C, cirrhosis, portal hypertension, esophageal varices, and thrombocytopenia 3 years ago. The treatment with steroids and interferon did not improve her condition. Mrs. M.P. was told there was a mass in her liver that was probably cancer. When she came to Dr. Berkson's office, she was fatigued, weak, pale, and her abdomen was enlarged. She was given alpha lipoic acid, silymarin, and selenium, along with the other nutrients. Her lab work began to show favorable changes, and she was able to recover her health and became free of the signs and symptoms of a serious chronic hepatitis C infection.

The second patient, Mrs. P.P., was a 49-year-old woman who was infected with hepatitis C more than 10 years ago following a blood transfusion before trauma surgery. A liver biopsy was taken about 3 years ago that showed moderate cirrhosis with active inflammation. She also developed portal hypertension with esophageal varices. She was treated with interferon therapy, but did not receive any results. A liver transplant was her only option and her health continued to deteriorate. She came to Dr. Berkson's office with symptoms of fatigue, anxiety, and insomnia. Mrs. P.P. was given 600 milligrams of alpha lipoic acid, 900 milligrams of silymarin, and 400 micrograms of selenium, along with aprazolam for the insomnia. She improved her diet and lifestyle and in 7 months regained her health. She was able to work a full time job and to play sports without fatigue or any other of her previous symptoms.

The third patient, Mrs. L.M., was a 35-year-old mother of 3 children. She developed hepatitis because of a blood transfusion she was given during the birth of her child 15 years ago. When she became ill three years ago, she was diagnosed with cirrhosis of the liver, portal hypertension, and esophageal varices. She also developed splenomegaly and thrombocytopenia as a result of her portal hypertension. It was recommended to Mrs. L.M. that she needed to get an evaluation for a liver transplant. When she came to Dr. Berkson's office, she complained of constant pain in the area of her liver and spleen. She was given the combination of alpha lipoic acid and other antioxidants. She began to feel much better within 2 weeks and recovered quickly. The pain in her liver ended and her energy level improved. She was able to return to her normal work as a housewife and returned to college the next semester earning a 3.8 grade point average.¹⁶⁴

Alpha lipoic acid has been shown to have a protective effect against acute hepatitis due to copper toxicity in experiments with rats. The Long-Evans Cinnamon rats accumulate excess copper in their liver due to a genetic defect. It causes them to develop acute hepatitis with severe jaundice in a manner similar to patients with Wilson's disease. Alpha lipoic acid was administered to the rats by gavage in doses of 10, 30, and 100 milligrams per kilogram of body weight five times per week. Alpha lipoic acid prevented the loss of body weight and the development of severe jaundice in a dose-dependant manner. Analysis of the antioxidant systems in the liver showed that alpha lipoic acid significantly suppressed the inactivations of catalase and glutathione peroxidase. It

completely suppressed the increase in lipid peroxidation and was found to exert antioxidant effects at the molecular level.¹⁶⁵

The protective effect of alpha lipoic acid on the liver has been studied in rats that suffered acute organ injuries. The rats were given a lipopolysaccharide to induce liver damage. The damage to the liver resulted in increases in plasma alanine levels, aspartate aminotransferase activities, tumor necrosis factor-alpha levels, and nitric oxide concentrations. There was also increased hepatic myeloperoxidase activity, increased lipid peroxidation, and increased levels of other markers that showed an activation of the inflammatory response. When the rats were pretreated with alpha lipoic acid prior to the liver injury, there was a general reversal of the altered biochemical indices and a significant alleviation of liver injuries.¹⁶⁶

Alpha lipoic acid may also benefit arthritis. In a study with mice, alpha lipoic acid was found to suppress the development of collagen-induced arthritis and to protect against bone destruction. The mice were divided into 3 groups and were given 10 milligrams per kilogram of body weight of alpha lipoic acid, 100 milligrams per kilogram of body weight of alpha lipoic acid, or a placebo. The alpha lipoic acid treated mice had a dose-dependant reduction of the collagen-induced arthritis. It also prevented bone erosion and destructive changes. Intracellular reactive oxygen species in lymphocytes obtained from inguinal lymph nodes was significantly reduced in the mice given alpha lipoic acid. TNF-alpha, IL-1beta, and IL-6, which are pro-inflammatory cytokines, were reduced by the treatment with alpha lipoic acid. The alpha lipoic acid was able to

improve the arthritis in the mice by the reduction of oxidative stress and the inhibiting of inflammatory cytokine activity.¹⁶⁷

There has been research on the treatment of human immunodeficiency virus (HIV) with alpha lipoic acid. Acquired immune deficiency syndrome (AIDS) is characterized by HIV infection and progressive depletion of T-cells and an impaired immune response. It is now widely accepted that one of the central features of HIV disease involves oxidative stress, which leads to cell death and a depletion of CD4 cells, or T-helper cells.¹⁶⁸ It has been demonstrated that the blood and tissues of HIV-infected individuals have increased oxidation reactions,¹⁶⁹ a deficiency of glutathione,¹⁷⁰ and increased amounts of free radicals.¹⁷¹ Glutathione is necessary for regulating immune system T-cell activation and phagocytosis.¹⁷² Alpha lipoic acid has the ability to raise glutathione levels as much as 30-70% both *in vivo* and *in vitro*, making it beneficial for HIV patients.¹⁷³ Raising glutathione levels has also been shown to alter the cytokine balance in favor of a Th1 immune response mode, an anticancer and antiviral mode of the immune system.¹⁷⁴ Patients with HIV have a compromised antioxidant defense system. Antioxidant levels in the blood are decreased and the peroxidation of lipids and proteins are typically increased in patients. In a pilot study, the short term effect of alpha lipoic acid on blood antioxidant levels and peroxidation products was investigated in HIV positive patients. In the majority of patients, alpha lipoic acid increased the plasma vitamin C levels, total glutathione levels, total blood thiol groups, and T helper lymphocyte levels, and decreased the lipid peroxidation products. The results of this

study show that alpha lipoic acid changes the blood redox state of HIV infected individuals.¹⁷⁵

Alpha lipoic acid has been found to be an effective inhibitor of HIV-1 virus replication. In cultured lymphoid T-cells, alpha lipoic acid was able to inhibit replication of the HIV virus and caused a 90% reduction in reverse transcriptase activity.¹⁷⁶ Nuclear factor-kappa beta (NF-kappaB) is one of the cellular transcription factors involved in gene expression of the human immunodeficiency virus. A randomized, double blind, placebo-controlled trial was done with 33 HIV-infected patients with a history of unresponsiveness to highly active antiretroviral treatment. They were given 900 milligrams of alpha lipoic acid per day or a placebo for 6 months. The glutathione level in the blood was significantly elevated after 6 months in the alpha lipoic acid-supplemented patients compared to insignificant changes in the patients taking the placebo. The lymphocyte proliferation response was significantly enhanced in the patients receiving alpha lipoic acid compared to the group taking a placebo. Alpha lipoic acid was found to have a positive impact on patients with HIV by restoring blood glutathione levels and improving the reactivity of lymphocytes to T-cell mitogens.¹⁷⁷

Because of its role as an antioxidant, alpha lipoic acid has been found to improve antibody response in immunosuppressed animals.¹⁷⁸ Alpha lipoic acid has been shown to cause cell death in some tumor cell lines.¹⁷⁹ Due to its ability to scavenge reactive oxygen species, alpha lipoic acid has been found to induce apoptosis, or cell death, in liver tumor cells. The apoptosis was dependant on the activation of the caspase cascade

and the mitochondrial death pathway. Alpha lipoic acid was able to increase caspase-9 and caspase-3 activity.¹⁸⁰ In a study using human cancer cells, alpha lipoic acid was found to activate an enzyme that kills leukemia cells. Leukemic Jurkat cells were treated with 100 microM of alpha lipoic acid for 72 hours. The enzyme that plays an important role in initiating cell death was increased 100% with alpha lipoic acid.¹⁸¹ Alpha lipoic acid was found to induce apoptosis in human colon cancer cells in another study. Alpha lipoic acid was found to increase caspase-3-like activity and increase the production of oxygen inside the mitochondria.¹⁸² Another study examined the effect of alpha lipoic acid on liver cancer cells. Two different hepatoma cell lines were used, FaO and HepG2. Alpha lipoic acid was shown to inhibit the growth of both cell lines as indicated by the reduction in the number of cells, the reduced expression of cyclin A, and the increased levels of the cyclin/CDKs inhibitors. This study showed that alpha lipoic acid induces apoptosis in liver cancer cells, and suggests that it may be useful in liver cancer therapy.¹⁸³

Alpha lipoic acid has been found to positively affect the long-term survival of a patient with pancreatic cancer that had metastasized to his liver. Following his diagnosis, he began a 21 day course of chemotherapy. Due to his poor tolerance for the chemotherapy, he decided to seek another opinion from a well-respected oncology center where he was told that there was little hope for his survival. When he came to Dr. Berkson's clinic, his quality of life was poor. He was losing weight and experiencing constant abdominal pain and nausea. He was put on intravenous alpha lipoic acid, 300-600 milligrams 2 days per week, and low-dose naltrexone, 4.5 milligrams at bedtime. In

addition, he was given an antioxidant regime of 600 milligrams oral alpha lipoic acid, 400 micrograms of selenium, and 300 milligrams 4 times a day of silymarin to reduce the products of oxidative stress that was due to his disorder. After the first intravenous administration of alpha lipoic acid, the patient commented that he had an increase of energy and a sense of well-being. His quality of life began to improve tremendously after starting the treatment. He went back to work soon after starting this therapy and remains free of symptoms 3 years later. There were no toxic adverse effects from the treatment and his malignancy has not progressed several years after the treatment.¹⁸⁴

Alpha lipoic acid may also be a valuable adjunct to chemotherapy. Studies with animals have shown that alpha lipoic acid is able to reduce the side effects associated with the chemotherapy drugs cyclophosphamide and vincristine, but did not hamper drug effectiveness.¹⁸⁵ Alpha lipoic acid has been shown to synergistically improve the cytotoxicity of vitamin C against cancer cells in tissue culture.¹⁸⁶ In another study using human tumor cell lines, alpha lipoic acid was shown to induce hyperacetylation of histones in vivo and shown to have a differential effect on the growth and viability of normal versus transformed cell lines. This study showed that human cancer cell lines became apoptotic after being exposed to alpha lipoic acid and that alpha lipoic acid did not cause apoptosis in normal cells.¹⁸⁷

Dihydro-lipoic acid has been found to inhibit skin tumor generation through its anti-inflammatory and antioxidant properties. Mice were given a topical application of 12-O-tetradecanoylphorbol on their skin. Dihydro-lipoic acid significantly inhibited the

stages of inflammation and inhibited tumor formation.¹⁸⁸ In a study with mice, alpha lipoic acid was investigated on a key inflammatory pathway and cell proliferation in cancerous ovarian epithelial cells. It was shown that alpha lipoic acid selectively inhibited the growth of the cancerous cells as compared to the non-cancerous cells of the ovary. The inhibitory effect was not due to apoptosis but instead is associated with an increase in the half-life of the cyclin-dependent kinase inhibitor, p27(kip1). The alpha lipoic acid also was found to inhibit TNF alpha-induced NF-kappa B signaling activity.¹⁸⁹ Rats with an aggressive cancer were treated with alpha lipoic acid in another study. The lifespan of the rats was increased by 25% with alpha lipoic acid.¹⁹⁰

In conclusion, in vitro and in vivo studies suggest that alpha lipoic acid acts as a powerful nutrient with diverse pharmacologic and antioxidant properties. Pharmacologically, alpha lipoic acid is able to improve glycemic issues, neuropathies, and protect against toxicities associated with heavy metal poisoning. As an antioxidant, alpha lipoic acid has a direct effect on eliminating free radicals, chelates metal ions, and increases glutathione and vitamin C levels. These diverse actions show how alpha lipoic acid can be effective in improving the pathology of many chronic diseases. Due to the fact that we are bombarded with a large amount of toxic chemicals in our daily environment,¹⁹¹ the human body is exposed to enormous amounts of chemicals that our liver must continually detoxify. This increases the amount of free radicals that we are exposed to, requiring the body to produce more antioxidants to neutralize the free radicals. The body's glutathione stores could be rapidly used up simply detoxifying the chemicals we are subjected to in our daily environment. This may be one of the reasons

for the decreased levels of glutathione in so many chronic diseases. From my research, oxidative stress and lowered glutathione levels are common factors that are involved in the pathology of many of our most common diseases. Because of its ability to improve glutathione levels, neutralize free radicals, and enhance liver function, alpha lipoic acid is one of the most beneficial supplements that people could take in order to help prevent and fight disease in the 21st century.

-
- ¹ Reed LJ. A trail of research from lipoic acid to a-keto acid dehydrogenase complexes. *J Biol Chem.* 2001 Oct 19;276(42):38329-36.
- ² Carreau JP. Biosynthesis of lipoic acid via unsaturated fatty acids. *Methods Enzymol.* 1979;62:152-158.
- ³ Bustamante J, Lodge JK, Marcocci L, Tritschler HJ, Packer L, Rihn BH. Alpha-lipoic acid in liver metabolism and disease. *Free Radic Biol Med.* 1998;24(6):1023-1039.
- ⁴ Morikawa T, Yasuno R, Wada H. Do mammalian cells synthesize lipoic acid? Identification of a mouse cDNA encoding a lipoic acid synthase located in mitochondria. *FEBS Lett.* 2001 Jun 1;498(1):16-21.
- ⁵ Harris RA, Joshi M, Jeoung NH, Obayashi M. Overview of the molecular and biochemical basis of branched-chain amino acid catabolism. *J Nutr.* 2005;135(6 Suppl):1527S-1530S.
- ⁶ Hofmann M, Mainka P, Tritschler H, Fuchs J, Zimmer G. Decrease of red cell membrane fluidity and –SH groups due to hyperglycemic conditions is counteracted by alpha-lipoic acid. *Arch Biochem Biophys.* 1995 Dec 1;324(1):85-92.
- ⁷ Biewenga GP, Haenen GR, Bast A. The pharmacology of the antioxidant lipoic acid. *Gen Pharmacol.* 1997 Oct;29(3):315-31.
- ⁸ Han D, Handelman G, Marcocci L, et al. Lipoic acid increases *de novo* synthesis of cellular glutathione by improving cystine utilization. *Biofactors.* 1997;6:321-338.
- ⁹ Hendler SS, Rorvik DR, eds. *PDR for Nutritional Supplements.* Montvale: Medical Economics Company, Inc; 2001.
- ¹⁰ Hermann R, Niebch G, Borbe HO, et al. Enantioselective pharmacokinetics and bioavailability of different racemic alpha-lipoic acid formulations in healthy volunteers. *Eur J Pharm Sci.* 1996;4:167-174.
- ¹¹ Gleiter CH, Schug BS, Hermann R, Elze M, Blume HH, Gundert-Remy U. Influence of food intake on the bioavailability of thioctic acid enantiomers. *Eur J Clin Pharmacol.* 1996;50(6):513-514.
- ¹² Bast A, Haenen GR. Interplay between lipoic acid and glutathione in the protection against microsomal lipid peroxidation. *Biochim Biophys Acta.* 1988 16;963(3):558-61.
- ¹³ Gurer H, et al. Antioxidant role of alpha lipoic acid in lead toxicity. *Free Radic Biol Med.* 1999 Jul 27(1-2):75-81.
- ¹⁴ Biewanga G, et al. The pharmacology of the antioxidant lipoic acid. *Gen Pharmac.* 1997 Sep;29(3):315-31.
- ¹⁵ Jones W, Li X, Qu ZC, Perriott L, Whitesell RR, May JM. Uptake, recycling, and antioxidant actions of alpha-lipoic acid in endothelial cells. *Free Radic Biol Med.* 2002 Jul 1;33(1):83-93.
- ¹⁶ Murray RK. Metabolism of xenobiotics. In: Murray RK, Granner DK, Mayes PA, et al, eds. *Harper's Biochemistry.* 24th ed. Stanford, CT: Appleton and Lange; 1996.
- ¹⁷ Hagen TM, Vinarsky V, Wehr CM, Ames BN. (R)-alpha-lipoic acid reverses the age-associated increase in susceptibility of hepatocytes to tert-butylhydroperoxide both in vitro and in vivo. *Antioxid Redox Signal.* 2000 Fall;2(3):473-483.
- ¹⁸ Suh JH, Shenvi SV, Dixon BM, et al. Decline in transcriptional activity of Nrf2 causes age-related loss of glutathione synthesis, which is reversible with lipoic acid. *Proc Natl Acad Sci U S A.* 2004 Mar 9;101(10):3381-3386.
- ¹⁹ Khanna S, Atalay M, Laaksonen DE, et al. Alpha-lipoic acid supplementation: tissue glutathione homeostasis at rest and after exercise. *J Appl Physiol.* 1999 Apr;86(4):1191-1196.
- ²⁰ Packer L, Tritschler HJ, Wessel K. Neuroprotection by the metabolic antioxidant alpha-lipoic acid. *Free Radic Biol Med.* 1997;22:359-378.
- ²¹ Han D, Handelman G, Marcocci L, et al. Lipoic acid increases *de novo* synthesis of cellular glutathione by improving cystine utilization. *Biofactors.* 1997;6(3):321-338.
- ²² Jain SK, Palmer M, Yanyun C. Effects of lipoic acid on glucose utilization, protein glycosylation and viscosity of high glucose-treated red blood cells. *Pathophysiology.* June 1998;5(1)75
- ²³ Thirunavukkarasu V, Anitha Nandhini AT, Anuradha CV. Lipoic acid improves glucose utilization and prevents protein glycation and AGE formation. *Pharmazie.* 2005 Oct;60(10):772-5.
- ²⁴ Midaoui AE, Elimadi A, Wu L, Haddad PS, de Champlain J. Lipoic acid prevents hypertension, hyperglycemia, and the increase in heart mitochondrial superoxide production. *Am J Hypertens.* 2003 Mar;16(3):173-9.
- ²⁵ Lynch MA. Lipoic acid confers protection against oxidative injury in non-neuronal and neuronal tissue. *Nutr Neurosci.* 2001;4(6):419-438.
- ²⁶ Gurer H, Ozgunes H, Oztezcan S, Ercal N. Antioxidant role of alpha-lipoic acid in lead toxicity. *Free Radic Biol Med.* 1999 Jul;27(1-2):75-81.

-
- ²⁷ Muller L, Menzel H. Studies on the efficacy of lipoate and dihydrolipoate in the alteration of cadmium²⁺ toxicity in isolated hepatocytes. *Biochem Biophys Acta*. 1990 May;1052(3):386-391.
- ²⁸ Sumathi R, Baskaran G, Varalakshmi P. Relationship between glutathione and DL alpha-lipoic acid against cadmium-induced hepatotoxicity. *Jap J Med Sci Biol*. 1996 Apr;49(2):39-48.
- ²⁹ Shila S, Kokilavani V, Subathra M, Panneerselvam C. Brain regional responses in antioxidant system to alpha-lipoic acid in arsenic intoxicated rat. *Toxicology*. 2005 May 15;210(1):25-36.
- ³⁰ Budhwar R, Kumar S. Prevention of chromate induced oxidative stress by alpha-lipoic acid. *Indian J Exp Biol*. 2005 Jun;43(6):531-5.
- ³¹ Packer L, Tritschler HJ, Wessel K. Neuroprotection by the metabolic antioxidant alpha-lipoic acid. *Free Radic Biol Med*. 1997;22(1-2): 359-78.
- ³² Rooney JP. The role of thiols, dithiols, nutritional factors and interacting ligands in the toxicology of mercury. *Toxicology*. 2007 May 20;234(3):145-156.
- ³³ Ramakrishnan N, Wolfe WW, Catravas GN. Radioprotection of hematopoietic tissues in mice by alpha lipoic acid. *Radiation Res*. 1992 Jun;130(3):360-5.
- ³⁴ Korkina L, et al. Antioxidant therapy in children affected by irradiation from the Chernobyl nuclear accident. *Biochem Soc Tran*. 1993 Aug;21(3):314S.
- ³⁵ Berkson BM. A conservative triple antioxidant approach to the treatment of hepatitis C. Combination of alpha lipoic acid (thioctic acid), silymarin, and selenium: three case histories. *Med Klin*. Oct 1999;94(3):84-9.
- ³⁶ Haramaki N, Assadnazari H, Zimmer G, et al. The influence of vitamin E and dihydrolipoic acid on cardiac energy and glutathione status under hypoxia-reoxygenation. *Biochem Mol Biol Int*. 1995 Oct;37(3):591-597.
- ³⁷ Yang R, Le G, Li A, Zheng J, Shi Y. Effect of antioxidant capacity on blood lipid metabolism and lipoprotein lipase activity of rats fed a high-fat diet. *Nutrition*. 2006 Nov-Dec;22(11-12):1185-1191.
- ³⁸ Yang RL, Li W, Shi YH, Le GW. Lipoic acid prevents high-fat diet-induced dyslipidemia and oxidative stress: A microarray analysis. *Nutrition*. 2008 Jun;24(6):582-8.
- ³⁹ Arivaxhagan P, Panneerselvam SR, Panneerselvam C. Effect of DL-alpha-lipoic acid on the status of lipid peroxidation and lipids in aged rats. *J Gerontol A Biol Sci Med Sci*. 2003 Sep;58(9):B788-91.
- ⁴⁰ Lenin M, Thiagarajan A, Nagaraj M, Varalakshmi P. Attenuation of oxalate-induced nephrotoxicity by eicosapentaenoate-lipoate (EPA-LA) derivative in experimental rat model. *Prostaglandins Leukot Essent Fatty Acids*. 2001 Nov-Dec;65(5-6):265-70.
- ⁴¹ Packer L. Alpha-Lipoic acid: a metabolic antioxidant which regulates NF-kappa B signal transduction and protects against oxidative injury. *Drug Metab Rev*. 1998;30(2):245-275.
- ⁴² Berkson BM, Rubin D, Berkson AJ. Long-term survival of a 46-year-old man with pancreatic cancer and liver metastases and treated with intravenous alpha lipoic acid and low dose naltrexone. *Integrative Cancer Therapies*. March 2006;5(1):83-89.
- ⁴³ Berkson BM, Rubin D, and Berkson AJ. Long term survival of a 46 year old man with pancreatic cancer and liver metastases and treated with intravenous alpha lipoic acid and low dose naltrexone. *Integrative Cancer Therapies*. 2006 March;5(1):83-89.
- ⁴⁴ Berkson BM, Rubin DM, Berkson AJ. Reversal of Signs and Symptoms of a B-cell lymphoma in a patient using only low-dose naltrexone *Integrative Cancer Therapies*. 2007 Sep;6(3):293-6.
- ⁴⁵ Henriksen EJ. Therapeutic effects of lipoic acid on hyperglycemia and insulin resistance. In: *Handbook of Antioxidants*, edited by Cadenas E and Packer L. New York: Dekker, 2001, p. 535-548.
- ⁴⁶ Evans JL, Goldfine ID. Alpha-lipoic acid: a multifunctional antioxidant that improves insulin sensitivity in patients with type 2 diabetes. *Diabetes Technol Ther*. 2000 Autumn;2(3):401-13.
- ⁴⁷ Packer L, et al. Molecular aspects of lipoic acid in the prevention of diabetes complications. *Nutrition*. 2001 Oct;17(10):888-95.
- ⁴⁸ Kamenova P. Improvement of insulin sensitivity in patients with type 2 diabetes mellitus after oral administration of alpha-lipoic acid. *Hormones*. 2006 Oct-Dec;5(4):251-258.
- ⁴⁹ Hagen TM, Yowe DL, Bartholomew JC, Wehr CM, Do KL, Park JY, Ames BN. Mitochondrial decay in hepatocytes from old rats: membrane potential declines, heterogeneity and oxidants increase. *Proc Natl Acad Sci USA*. 1997 Apr 1;94(7):3064-3069.
- ⁵⁰ Hagen TM, Ingersoll RT, et al. R-lipoic acid-supplemented old rats have improved mitochondrial function, decreased oxidative damage, and increased metabolic rate. *FASEB J*. 1999 Feb;13(2):411-418.

-
- ⁵¹ Ha H, Lee JH, Kim HN, et al. Alpha-lipoic acid inhibits inflammatory bone resorption by suppressing prostaglandin E2 synthesis. *J Immunol.* 2006 Jan 1;176(1):111-7.
- ⁵² Jesudason EP, Masilamoni JG, Jebaraj CE, Paul SF, Jayakumar R. Efficacy of DL-alpha lipoic acid against systemic inflammation-induced mice: antioxidant defense system. *Mol Cell Biochem.* 2008 Jun;313(1-2):113-23.
- ⁵³ Masood A, Banerjee B, Vijayan VK, et al. Modulation of stress-induced neurobehavioral changes by nitric oxide in rats. *Eur J Pharmacol.* 2003 Jan 1;458(1-2):135-9.
- ⁵⁴ Olivenza R, Moro MA, Lizasoain I, et al. Chronic stress induces the expression of inducible nitric oxide synthase in rat brain cortex. *J Neurochem.* 2000 Feb;74(2):785-91.
- ⁵⁵ Zaidi R, et al. Modulation of restraint stress induced oxidative changes in rats by antioxidant vitamins. *Journal of Nutritional Biochemistry.* 2003 Nov;14(11):633-36.
- ⁵⁶ Akpınar D, Yargıçoğlu P, Derin N, Aliciğuzel Y, Agar A. The effect of lipoic acid on antioxidant status and lipid peroxidation in rats exposed to chronic restraint stress. *Physiol Res.* 2007 Nov 30.
- ⁵⁷ Ohmori H, Yamauchi T, Yamamoto I. Augmentation of the antibody response by lipoic acid in mice. II. Restoration of the antibody response in immunosuppressed mice. *Jpn J Pharmacol.* 1986 Oct;42(2):275-280.
- ⁵⁸ Fomichev VI, Pchelintsev VP. The neurohumoral systems of patients with ischemic heart disease and under emotional pain-stress: the means for their pharmacological regulation. *Kardiologıia.* 1993;33(10):15-18.
- ⁵⁹ Rao GN, Sadasivudu B, Cotlier E. Studies of glutathione s-transferase, glutathione peroxidase and glutathione reductase in human normal and cataractous lenses. *Ophthalmic Res.* 1983;15(4):173-179.
- ⁶⁰ Maitra I, et al. Alpha-lipoic acid prevents buthionine sulfoximine-induced cataract formation in newborn rats. *Free Rad Biol Med.* 1995 Apr;18(4):823-9.
- ⁶¹ Maitra I, Serbinova E, Tritschler HJ, Packer L. Stereospecific effects of R-lipoic acid on buthionine sulfoximine-induced cataract formation in newborn rats. *Biochem Biophys Res Commun.* 1996 Apr;221(2):422-29.
- ⁶² Filina AA, Davydova NG, Kolomoitseva EM. The effect of lipoic acid on the components of the glutathione system in the lacrimal fluid of patients with open-angle glaucoma. *Vestn Oftalmol.* 1993;109:5-7. [Article in Russian]
- ⁶³ Filina AA, Davydova NG, Endrikhovskii SN, Shamshinova AM. Lipoic acid as a means of metabolic therapy of open-angle glaucoma. *Vestn Oftalmol.* 1995;1 1:6-8. [Article in Russian]
- ⁶⁴ Shigenaga MK, Hagen TM, Ames BN. Oxidative damage and mitochondrial decay in aging. *Proc Natl Acad Sci USA.* 1994 Nov 8;91(23):10771-10778
- ⁶⁵ Hagen TM, Yowe DL, Bartholomew JC, Wehr CM, Do KL, Park JY, Ames BN. Mitochondrial decay in hepatocytes from old rats: membrane potential declines, heterogeneity and oxidants increase. *Proc Natl Acad Sci USA.* 1997 Apr 1;94(7):3064-3069
- ⁶⁶ Ames BN, Shigenaga MK, Hagen TM. Oxidants, antioxidants, and the degenerative diseases of aging. *Proc Natl Acad Sci USA.* 1993 Sep 1;90(17):7915-7922.
- ⁶⁷ Hagen TM, Ingersoll RT, Lykkesfeldt J, et al. (R)-alpha-lipoic acid-supplemented old rats have improved mitochondrial function, decreased oxidative damage, and increased metabolic rate. *FASEB J.* 1999 Feb;13(2):411-8.
- ⁶⁸ Hagen TM, Liu J, Lykkesfeldt J, Wehr CM, et al. Feeding acetyl-L-carnitine and lipoic acid to old rats significantly improves metabolic function while decreasing oxidative stress. *Proc Natl Acad Sci USA.* 2002 Feb 19;99(4):1870-5.
- ⁶⁹ Beutner H. Randomized, placebo-controlled, double blind study on the clinical efficacy of a cream containing 5% alpha-lipoic acid related to photoageing of the facial skin. *Br J Dermatol.* 2003;149:841-9.
- ⁷⁰ Liu J, Atamna H, Kuratsune H, Ames BN. Delaying brain mitochondrial decay and aging with mitochondrial antioxidants and metabolites. *Ann N Y Acad Sci.* 2002 May;959:133-66.
- ⁷¹ Liu J, Killilea DW, Ames BN. Age-associated mitochondrial oxidative decay: improvement of carnitine acetyltransferase substrate-binding affinity and activity in brain by feeding old rats acetyl-L-carnitine and/or R-alpha lipoic acid. *Proc Natl Acad Sci USA.* 2002 Mar 19;99(4):1876-81.
- ⁷² Milgram NW, Araujo JA, Hagen TM, et al. Acetyl-L-carnitine and alpha-lipoic acid supplementation of aged beagle dogs improves learning in two landmark discrimination tests. *FASEB J.* 2007 Nov;21(13):3756-62.

-
- ⁷³ Holmquist L, Stauchbury G, et al. Lipoic acid as a novel treatment for Alzheimer's disease and related dementias. *Pharmacol Ther.* 2007 Jan;113(1):154-64.
- ⁷⁴ Hager K, Marahrens A, Kenklies M, Riederer P, Munch G. Alpha-lipoic acid as a new treatment option for Alzheimer type dementia. *Arch Gerontol Geriatr.* 2001 Jul;32(3):275-82.
- ⁷⁵ Lovell MA, Xie C, Xiong S, Markesbery WR. Protection against amyloid beta peptide and iron/hydrogen peroxide toxicity by alpha lipoic acid. *J Alzheimers Dis.* 2003 Jul;5(3):229-39.
- ⁷⁶ Karunakaran S, Diwakar L, Saeed U, et al. Activation of apoptosis signal regulating kinase 1 (ASK1) and translocation of death-associated protein, Daxx, in substantia nigra pars compacta in a mouse model of Parkinson's disease: protection by alpha-lipoic acid. *FASEB J.* 2007 Jul;21(9):2226-36.
- ⁷⁷ Pershandsingh HA. Alpha-lipoic acid: physiologic mechanisms and indications for the treatment of metabolic syndrome. *Expert Opin Investig Drugs.* 2007 Mar;16(3):291-302.
- ⁷⁸ Kim MS, Park JY, Namkoong C, et al. Anti-obesity effects of alpha-lipoic acid mediated by suppression of hypothalamic AMP-activated protein kinase. *Nat Med.* 2004 Jul;10(7):727-33.
- ⁷⁹ Sola S, Mir MQ, Cheema FA, et al. Irbesartan and lipoic acid improve endothelial function and reduce markers of inflammation in the metabolic syndrome: results of the Irbesartan and Lipoic Acid in Endothelial Dysfunction (ISLAND) study. *Circulation.* 2005 Jan 25;111(3):343-8.
- ⁸⁰ Heller B, Burkhardt V, Lampeter E, Kolb H. Antioxidant therapy for the prevention of type I diabetes. *Adv Pharmacol.* 1997;38:629-638.
- ⁸¹ Hagen TM, et al. Feeding acetyl-L-carnitine and lipoic acid to old rats significantly improves the metabolic function while decreasing oxidative stress. *Proc Natl Acad Sci USA.* 2002 Feb 19;99(4):1870-5.
- ⁸² Evans JL, Goldfine ID. Alpha-lipoic acid: a multifunctional antioxidant that improves insulin sensitivity in patients with type 2 diabetes. *Diabetes Technol Ther.* 2000 Autumn;2(3):401-13.
- ⁸³ Jacob S, Henriksen EJ, Tritschler HJ, Augustin HJ, Dietze GJ. Improvement of insulin-stimulated glucose-disposal in type 2 diabetes after repeated parenteral administration of thioctic acid. *Exp Clin Endocrinol Diabetes.* 1996;104(3):284-8.
- ⁸⁴ Jacob S, Henriksen EJ, Schiemann AL, et al. Enhancement of glucose disposal in patients with type 2 diabetes by alpha-lipoic acid. *Arzneimittelforschung.* 1995;45(8):872-874.
- ⁸⁵ Jacob S, Rett K, Henriksen EJ, Haring HU. Thioctic acid--effects on insulin sensitivity and glucose-metabolism. *Biofactors.* 1999;10(2-3):169-174.
- ⁸⁶ Ziegler D. Thioctic acid for patients with symptomatic diabetic polyneuropathy: a critical review. *Treat Endocrinol.* 2004;3(3):173-189.
- ⁸⁷ Konrad T, Vicini P, Kusterer K, et al. Alpha lipoic acid treatment decreases serum lactate and pyruvate concentrations and improves glucose effectiveness in lean and obese patients with type 2 diabetes. *Diabetes Care.* 1999 Feb;22(2):280-287.
- ⁸⁸ Streeper RS, Henriksen EJ, Jacob S, et al. Differential effects of lipoic acid stereoisomers on glucose metabolism in insulin-resistant skeletal muscle. *Am J Physiol.* 1997 Jul;273(1 Pt 1):E185-E191.
- ⁸⁹ Estrada DE, Ewart HS, Tsakiridis T, et al. Stimulation of glucose uptake by the natural coenzyme alpha-lipoic acid/thioctic acid: participation of elements of the insulin signaling pathway. *Diabetes.* 1996 Dec;45(12):1798-1804.
- ⁹⁰ Kramer K, Packer L. R-alpha-lipoic acid. In: Kramer K, Hoppe P, Packer L, eds. *Nutraceuticals in Health and Disease Prevention.* New York: Marcel Dekker, Inc.; 2001:129-164.
- ⁹¹ Ziegler D, Hanefeld M, Ruhnau KJ, et al. Treatment of symptomatic diabetic peripheral neuropathy with the anti-oxidant alpha-lipoic acid. A 3-week multicentre randomized controlled trial (ALADIN study). *Diabetologia.* 1995 Dec;38(12):1425-1433.
- ⁹² Reljanovic M, Reichel G, Rett K, et al. Treatment of diabetic polyneuropathy with the antioxidant thioctic acid (alpha-lipoic acid): a two year multicenter randomized double-blind placebo-controlled trial (ALADIN II). Alpha Lipoic Acid in Diabetic Neuropathy. *Free Radic Res.* 1999 Sept;31(3):171-179.
- ⁹³ Ziegler D, Hanefeld M, Ruhnau KJ, et al. Treatment of symptomatic diabetic polyneuropathy with the antioxidant alpha-lipoic acid: a 7-month multicenter randomized controlled trial (ALADIN III Study). ALADIN III Study Group. Alpha-Lipoic Acid in Diabetic Neuropathy. *Diabetes Care* 1999 Aug;22(8):1296-1301.
- ⁹⁴ Ametov AS, Barinov A, Dyck PJ, et al. The sensory symptoms of diabetic polyneuropathy are improved with alpha-lipoic acid: the AYDNEY trial. *Diabetes Care.* 2003;26:770-776.
- ⁹⁵ Ziegler D, Nowak H, Kempler P, Vargha P, Low PA. Treatment of symptomatic diabetic polyneuropathy with the antioxidant alpha-lipoic acid: a meta-analysis. *Diabet Med.* 2004;21(2):114-121.

-
- ⁹⁶ Ruhnau KJ, Meissner HP, Finn JR, et al. Effects of 3-week oral treatment with the antioxidant thioctic acid in symptomatic diabetic polyneuropathy. *Diabet Med.* 1999 Dec;16(12):1040-1043.
- ⁹⁷ Reljanovic M, Reichel G, Rett K, et al. Treatment of diabetic polyneuropathy with the antioxidant thioctic acid (alpha-lipoic acid): a two year multicenter randomized double-blind placebo-controlled trial (ALADIN II). *Alpha Lipoic Acid in Diabetic Neuropathy. Free Radic Res.* 1999;31(3):171-179.
- ⁹⁸ Ziegler D. Thioctic acid for patients with symptomatic diabetic polyneuropathy: a critical review. *Treat Endocrinol.* 2004;3(3):173-189.
- ⁹⁹ Ziegler D, Schatz H, Conrad F, Gries FA, Ulrich H, Reichel G. Effects of treatment with the antioxidant alpha-lipoic acid on cardiac autonomic neuropathy in NIDDM patients. A 4-month randomized controlled multicenter trial (DEKAN Study). *Deutsche Kardiale Autonome Neuropathie. Diabetes Care.* 1997;20(3):369-373.
- ¹⁰⁰ Yi X and Maeda N. Alpha-Lipoic Acid Prevents the Increases in Atherosclerosis Induced by Diabetes in Apolipoprotein E-Deficient Mice Fed High Fat/Low Cholesterol Diet. *Diabetes.* 55(8):2238-44 (2006).
- ¹⁰¹ Schalkwijk CG, Stehouwer CD. Vascular complications in diabetes mellitus: the role of endothelial dysfunction. *Clin Sci. (Lond).* 2005;109(2):143-159.
- ¹⁰² Sola S, Mir MQ, Cheema FA, et al. Irbesartan and lipoic acid improve endothelial function and reduce markers of inflammation in the metabolic syndrome: results of the Irbesartan and Lipoic Acid in Endothelial Dysfunction (ISLAND) study. *Circulation.* 2005;111(3):343-348.
- ¹⁰³ Lateef H, Aslam MN, Stevens MJ, Varani J. Pretreatment of diabetic rats with lipoic acid improves healing of subsequently-induced abrasion wounds. *Arch Dermatol Res.* 2005 Aug;297(2):75-83.
- ¹⁰⁴ Alleva R, Tomasetti M, Sartini D, Emanuelli M, et al. Alpha-lipoic acid modulates extracellular matrix and angiogenesis gene expression in non-healing wounds treated with hyperbaric oxygen therapy. *Mol Med.* 2008 Mar-Apr;14(3-4):175-83.
- ¹⁰⁵ Schreibelt G, Musters RJ, Reijerkerk A, et al. Lipoic acid affects cellular migration into the central nervous system and stabilizes blood-brain barrier integrity. *J Immunol.* 2006 Aug 15;177(4):2630-7.
- ¹⁰⁶ Marracci GH, Jones RE, McKeon GP, Bourdette DN. Alpha lipoic acid inhibits T cell migration into the spinal cord and suppresses and treats experimental autoimmune encephalomyelitis. *J Neuroimmunol.* 2002;131(1-2):104-114.
- ¹⁰⁷ Yadav V, Marracci G, Lovera J, et al. Lipoic acid in multiple sclerosis: a pilot study. *Mult Scler.* 2005;11(2):159-165.
- ¹⁰⁸ Magis D, Ambrosini A, Sandor P, et al. A randomized double-blind placebo-controlled trial of thioctic acid in migraine prophylaxis. *Headache.* 2007 Jan;47(1):52-7.
- ¹⁰⁹ Hoffman MA, Tritschler HJ, Bierhaus A, Zeigler R, Wahl P, Nawroth P. Lipoate effects on atherogenesis. Fuchs, J. Packer, L. Zimmer, G. eds. *Lipoic Acid in Health and Disease.* 2000:321-336 Marcel Dekker New York. .
- ¹¹⁰ Tardif JC, Bourassa MG. eds. *Developments in Cardiovascular Medicine: Antioxidants and Cardiovascular Disease.* v. 233 2000 Kluwer Academic Press The Netherlands. .
- ¹¹¹ Wollin SD, and Jones PH. Alpha-lipoic acid and Cardiovascular disease. *J. Nutr.* 2003 Nov; 133(11): 3327-3330.
- ¹¹² Shih JCH. Atherosclerosis in Japanese quail and the effect of lipoic acid. *Fed Proc.* 1983 May;42(8):2494-7.
- ¹¹³ Ivanov VN. Effect of lipoic acid on tissue respiration in rabbits with experimental atherosclerosis. *Cor Vasa.* 1974;16(2):141-150.
- ¹¹⁴ Zhang WJ, Bird KE, McMillen TS, Leboeuf RC, et al. Dietary alpha-lipoic acid supplementation inhibits atherosclerotic lesion development in apolipoprotein E deficient and apolipoprotein E/low-density lipoprotein receptor deficient mice. *Circulation.* 2008 Jan 22;117(3): 421-428.
- ¹¹⁵ Gao TL, Huang YZ. Effects of lipoic acid on reperfusion induced arrhythmias and myocardial action potential alterations induced by free radical generating system. *Sheng Li Xue Bao.* 1991 Apr;43(2):149-55.
- ¹¹⁶ Kumar KV, Das UN. Are free radicals involved in the pathobiology of human essential hypertension? *Free Radic Res Commun.* 1993;19(1):59-66.
- ¹¹⁷ Vasdev S, Ford CA, Parai S, Longerich L, Gadag V. Dietary alpha-lipoic acid supplementation lowers blood pressure in spontaneously hypertensive rats. *J Hypertens.* 2000 May;18(5):567-73.
- ¹¹⁸ El Midaoui A, de Champlain J. Prevention of hypertension, insulin resistance, and oxidative stress by alpha-lipoic acid. *Hypertension.* 2002 Feb;39(2):303-7.

-
- ¹¹⁹ Kim MS, Park JY, Namkoong C, Jang PG, Ryu JW, Song HS, Yun JY, Namgoong IS, Ha J, Park IS, Lee IK, Viollet B, Youn JH, Lee HK, Lee KU. Anti-obesity effects of alpha-lipoic acid mediated by suppression of hypothalamic AMP-activated protein kinase. *Nat Med*. 2004 Jul; 10(7): 727–733.
- ¹²⁰ Lee WJ, Song KH, Koh EH, Won JC, Kim HS, Park HS, Kim MS, Kim SW, Lee KU, Park JY. alpha-Lipoic acid increases insulin sensitivity by activating AMPK in skeletal muscle. *Biochem Biophys Res Commun*. 2005 Jul 8; 332(3): 885–891.
- ¹²¹ Poulter N. Global risk of cardiovascular disease. *Heart*. 2003; 89 (suppl 2): ii2–ii5.
- ¹²² Sakamoto S, Minami K, Niwa Y, Ohnaka M, Nakaya Y, Mizuno A, Kuwajima M, Shima K. Effect of exercise training and food restriction on endothelium-dependent relaxation in the Otsuka Long-Evans Tokushima Fatty rat, a model of spontaneous NIDDM. *Diabetes*. 1998 Jan; 47(1): 82–86.
- ¹²³ Ross R. Atherosclerosis—an inflammatory disease. *N Engl J Med*. 1999; 340: 115–126.
- ¹²⁴ Lee WJ, Lee IK, Kim HS, et al. Alpha-lipoic acid prevents endothelial dysfunction in obese rats via activation of AMP-activated protein kinase. *Arterioscler Thromb Vasc Biol*. 2005 Dec;25(12):2488-94.
- ¹²⁵ Mitsui Y, Schmelzer JD, Zollman PJ, et al. Alpha-lipoic acid provides neuroprotection from ischemia-reperfusion injury of peripheral nerve. *J Neurol Sci*. 1999 Feb 1;163(1):11-16.
- ¹²⁶ Haramaki N, Assadnazari H, Zimmer G, et al. The influence of vitamin E and dihydrolipoic acid on cardiac energy and glutathione status under hypoxia-reoxygenation. *Biochem Mol Biol Int*. 1995 Oct;37(3):591-597.
- ¹²⁷ Cao X, Phillis JW. The free radical scavenger, alpha-lipoic acid, protects against cerebral ischemia-reperfusion injury in gerbils. *Free Rad Res*. 1995 Oct;23(4):365-370.
- ¹²⁸ Panigrahi M, Sadguna Y, Shivakumar BR, et al. Alpha-lipoic acid protects against reperfusion injury following cerebral ischemia in rats. *Brain Res*. 1996 Apr 22;717(1-2):184-188.
- ¹²⁹ Serbinova E, Khwaja S, Reznick AA, Packer L. Thiocetic acid protects against ischemia-reperfusion injury in the isolated perfused Langendorff heart. *Free Rad. Res Comm*. 1992;17(1):49-58.
- ¹³⁰ Clark W, Rinker L, Lessov N, Lowery S, Cippola M. Efficacy of antioxidant therapies in transient focal ischemia in mice. *Stroke*. 2001 Apr;32(4):1000-4.
- ¹³¹ Fowler BA. General subcellular effects of lead, mercury, cadmium, and arsenic. *Environ Health Perspect*. 1978 Feb;22:37-41.
- ¹³² Grunert RR. The effect of dl-alpha-lipoic acid on heavy metal intoxication in mice and dogs. *Arch Biochem Biophys*. 1960 Feb;86:190-4.
- ¹³³ Miura K, Clarkson TW. Reduced methylmercury accumulation in a methylmercury resistant rat pheochromocytoma PC12 cell line. *Toxicol Appl Pharmacol*. 1993 Jan;118(1):39-45.
- ¹³⁴ Bustamante J, Lodge JK, Marcocci L, et al. Alpha-Lipoic acid in liver metabolism and disease. *Free Radic Biol Med*. 1998 Apr;24(6):1023-1039.
- ¹³⁵ Muller L, Menzel H. Studies on the efficacy of lipoate and dihydrolipoate in the alteration of [cadmium²⁺] toxicity in isolated hepatocytes. *Biochim Biophys Acta*. 1990 May 22;1052(3):386-391.
- ¹³⁶ Sumathi R, Devi VK, Varalakshmi P. DL-alpha lipoic acid protection against cadmium-induced tissue lipid peroxidation. *Med Sci Res*. 1994;22: 23-25.
- ¹³⁷ Lodge JK, Traber MG, Packer L. Thiol chelation of [Cu²⁺] by dihydrolipoic acid prevents human low density lipoprotein peroxidation. *Free Radic Biol Med*. 1998 Aug;25(3):287-297.
- ¹³⁸ de Costa VS, Morbus DA. Wilson's disease and the possible therapeutic role of alpha lipoic acid. *Arzneimittelforschung*. 1970;20:1210-1213.
- ¹³⁹ Packer L, Tritschler HJ, Wessel K. Neuroprotection by the metabolic antioxidant alpha-lipoic acid. *Free Radio Biol Med*. 1997;22(1-2):359-378.
- ¹⁴⁰ Somani SM, Husain K, Whitworth C, et al. Dose-dependent protection by lipoic acid against cisplatin-induced nephrotoxicity in rats: antioxidant defense system. *Pharmacol Toxicol*. 2000 May;86(5):234-241.
- ¹⁴¹ Gurer H, Ozgunes H, Oztezcan S, Ercal N. Antioxidant role of alpha-lipoic acid in lead toxicity. *Free Radic Biol Med*. 1999 Jul;27(1-2):75-81.
- ¹⁴² Femiano F, Scully C, Gombos F. Idiopathic dysgeusia; an open trial of alpha lipoic acid (ALA) therapy. *Int J Oral Maxillofac Surg*. 2002 Dec;31(6): 625-628.
- ¹⁴³ Femiano F, Scully C. Burning mouth syndrome (BMS):double blind controlled study of alpha-lipoic acid (thioctic acid) therapy. *J Oral Pathol Med*. 2002 May; 31(5): 267-269.
- ¹⁴⁴ Byun CH, Koh JM, Kim DK, Park SI, Lee KU, Kim GS. Alpha-lipoic acid inhibits TNF-alpha-induced apoptosis in human bone marrow stromal cells. *J Bone Miner Res*. 2005 Jul;20(7):1125-35.

-
- ¹⁴⁵ Kim HJ, Chang EJ, Kim HM, Lee SB, et al. Antioxidant alpha-lipoic acid inhibits osteoclast differentiation by reducing nuclear factor-kappaB DNA binding and prevents in vivo bone resorption induced by receptor activator of nuclear factor-kappaB ligand and tumor necrosis factor-alpha. *Free Radic Biol Med*. 2006 May 1;40(9):1483-93.
- ¹⁴⁶ Seidman MD, Khan MJ, Bai U, Shirwany N, Quirk WS. Biologic activity of mitochondrial metabolites on aging and age-related hearing loss. *Am J Otol*. 2000 Mar;21(2):161-167.
- ¹⁴⁷ Rybak L, Husain K, Morris C, Whitworth C, Somani S. Effect of protective agents against cisplatin ototoxicity. *Am J Otol*. 2000 Jul; 21(4):513-520.
- ¹⁴⁸ Rybak LP, Husain K, Whitworth C, Somani SM. Dose dependent protection by lipoic acid against cisplatin-induced ototoxicity in rats: antioxidant defense system. *Toxicol Sci*. 1999 Feb;47(2):195-202.
- ¹⁴⁹ Conlon BJ, Aran JM, Erre JP, Smith DW. Attenuation of aminoglycoside-induced cochlear damage with the metabolic antioxidant alpha-lipoic acid. *Hear Res* 1999 Feb;128(1-2):40-4.
- ¹⁵⁰ Armstrong KL, Cooper MF, Williams MT, Elsayed NM. Vitamin E and lipoic acid, but not vitamin C improve blood oxygenation after high-energy IMPULSE noise (BLAST) exposure. *Biochem Biophys Res Commun*. 1998 Dec 9;253(1):114-8.
- ¹⁵¹ Prahalathan C, Selvakumar E, Varalakshmi P. Lipoic acid modulates adriamycin-induced testicular toxicity. *Reprod Toxicol*. 2006 Jan;21(1):54-9.
- ¹⁵² Prahalathan C, Selvakumar E, Varalakshmi P. Lipoic acid ameliorates adriamycin-induced testicular mitochondriopathy. *Reprod Toxicol*. 2005 May-Jun;20(1):111-6.
- ¹⁵³ Malarkodi KP, Sivaprasad R, Varalakshmi P. Effect of lipoic acid on the oxidoreductive status of red blood cells in rats subject to oxidative stress by chronic administration of adriamycin. *Hum Exp Toxicol*. 2004 Mar;23(3):129-35.
- ¹⁵⁴ Abdel-Zaher AO, Abdel-Hady RH, Mahmoud MM, Farraq MM. The potential protective role of alpha-lipoic acid against acetaminophen-induced hepatic and renal damage. *Toxicology*. 2008 Jan 20;243(3):261-70.
- ¹⁵⁵ Makeeva AV, Popova TN, Matasova LV. [Thioctic acid action on glutathione-dependant antioxidant system functioning at toxic hepatitis of rats] *Biomed Khim*. 2007 Mar-Apr;53(2):181-9.
- ¹⁵⁶ Moller E, Schmitt R. [A contribution to the treatment of chronic liver diseases (author's tranl)] *Med Klin*. 1976 Oct 22;71(43):1831-5.
- ¹⁵⁷ Lykkesfeldt J, Christen S, Wallock LM, Chang HH, Jacob RA, Ames BN. Ascorbate is depleted by smoking and repleted by moderate supplementation: a study in male smokers and nonsmokers with matched dietary antioxidant intakes. *Am J Clin Nutr*. 2000 Feb;71(2):530-536.
- ¹⁵⁸ Cross CE, O'Neill CA, Reznick AZ, et al. Cigarette smoke oxidation of human plasma constituents. *Ann NY Acad Sci*. 1993 May 28;686:72-89.
- ¹⁵⁹ Tomany SC, Wang JJ, Van Leeuwen R, et al. Risk factors for incident age-related macular degeneration: pooled findings from 3 continents. *Ophthalmology*. 2004 Jul;111(7):1280-1287
- ¹⁶⁰ Eiserich JP, van der Vliet A, Handelman GJ, Halliwell B, Cross CE. Dietary antioxidants and cigarette smoke-induced biomolecular damage: a complex interaction. *Am J Clin Nutr*. 1995;62(suppl 6)1490S-1500S.
- ¹⁶¹ Esterbauer H, Schaur RJ, Zollner H. Chemistry and biochemistry of 4-hydroxynonenal, malonaldehyde and related aldehydes. *Free Radic Biol Med*. 1991;11(1):81-128.
- ¹⁶² Jia L, Liu Z, Sun L, Miller SS, et al. Acrolein, a toxicant in cigarette smoke, causes oxidative damage and mitochondrial dysfunction in RPE cells: protection by (R)-alpha-lipoic acid. *Invest Ophthalmol Vis Sci*. 2007 Jan;48(1):339-48.
- ¹⁶³ Berkson, BM. "Thioctic Acid in the Treatment of Poisoning with Alpha amanitin." *Amanita Toxins and Poisonings*, 1980. *Amanita Toxins and Poisonings*, 203 (Heidelberg: International Amanita Symposium, Nov. 1-3, 1978). eds Faulstich, H., Kommerell, B., and Th. Wieland, Verlag Gerhard Witzstrock, Baden-Baden, Koln, New York 1980.
- ¹⁶⁴ Berkson BM. A conservative triple antioxidant approach to the treatment of hepatitis C. Combination of alpha lipoic acid (thioctic acid), silymarin, and selenium: three case histories. *Med Klin*. (Munich) 1999 Oct 15;94 Suppl 3:84-9.
- ¹⁶⁵ Yamamoto H, Watanabe T, Mizuno H, et al. The antioxidant effect of DL-alpha-lipoic acid on copper-induced acute hepatitis in Long-Evans Cinnamon (LEC) rats. *Free Radic Res*. 2001 Jan;34(1):69-80.
- ¹⁶⁶ Suntres ZE. Prophylaxis against lipopolysaccharide-induced liver injuries by lipoic acid in rats. *Pharmacol Res*. 2003 Dec;48(6):585-91.

- ¹⁶⁷ Lee EY, Lee CK, Lee KU, Park JY, et al. Alpha-lipoic acid suppresses the development of collagen-induced arthritis and protects against bone destruction in mice. *Rheumatol Int.* 2007 Jan;27(3):225-33.
- ¹⁶⁸ Pace GW, Leaf CD. The role of oxidative stress in HIV disease. *Free Radic Biol Med.* 1995 Oct;19(4):523-528.
- ¹⁶⁹ Sato N, Iwata S, Nakamura K, et al. Thiol-mediated redox regulation of apoptosis. Possible roles of cellular thiols other than glutathione in T cell apoptosis. *J Immunol.* 1995 Apr 1;154(7):3194-3203.
- ¹⁷⁰ Staal FJ, Roederer M, Israelski DM, et al. Intracellular glutathione levels in T cell subsets decrease in HIV-infected individuals. *AIDS Res Hum Retroviruses.* 1992 Feb;8(2):305-311.
- ¹⁷¹ Greenspan HC, Aruoma OI. Oxidative stress and apoptosis in HIV infection: a role for plant-derived metabolites with synergistic antioxidant activity. *Immunol Today.* 1994 May;15(5):209-213.
- ¹⁷² Hamilos DL, Wedner HJ. The role of glutathione in lymphocyte activation. I. Comparison of inhibitory effects of buthionine sulfoximine and 2-cyclohexene-1-one by nuclear size transformation. *J Immunol.* 1985 Oct;135(4):2740-2747.
- ¹⁷³ Han D, Tritschler HJ, Packer L. Alpha-lipoic acid increases intracellular glutathione in a human T-lymphocyte Jurkat cell line. *Biochem Biophys Res Commun.* 1995 Feb 6;207(1):258-264.
- ¹⁷⁴ Peterson JD, Herzenberg LA, Vasquez K, Waltenbaugh C. Glutathione levels in antigen-presenting cells modulate Th1 versus Th2 response patterns. *Proc Natl Acad Sci USA.* 1998 Mar 17;95(6):3071-6.
- ¹⁷⁵ Fuchs J, Schofer H, Milbradt R, et al. Studies on lipoate effects on blood redox state in human immunodeficiency virus infected patients. *Arzneimittelforschung.* 1993 Dec;43(12):1359-62.
- ¹⁷⁶ Baur A, Harrer T, Peukert M, et al. Alpha-lipoic acid is an effective inhibitor of human immunodeficiency virus (HIV-1) replication. *Klin Wochenschr.* 1991 Oct 2;69(15):722-4.
- ¹⁷⁷ Jariwalla RJ, Lalezari J, Cenko D, Mansour SE, Kumar A, et al. Restoration of blood glutathione status and lymphocyte function following alpha-lipoic acid supplementation in patients with HIV infection. *J Altern Complement Med.* 2008 Mar;14(2):139-46.
- ¹⁷⁸ Ohmori Hnt, Yamauchi T, Yamamoto I. Augmentation of the antibody response by lipoic acid in mice. 1. Analysis of the mode of action in an in vitro cultures system. *Jpn J Pharmacol.* 1986 Sep;42(1):135-40.
- ¹⁷⁹ Van de MK, Chen JS, Steliou K, Perrine SP, Fatter DV. Alpha-lipoic acid induces p27Kip-dependant cell cycle arrest in non-transformed cell lines and apoptosis in tumor cell lines. *J Cell Physiol.* 2003 Mar;194(3):325-40.
- ¹⁸⁰ Shi DY, Liu HL, Stern JS, Yu PZ, Liu SL. Alpha-lipoic acid induces apoptosis in hepatoma cells via the PTEN/Akt pathway. *FEBS Lett.* 2008 May 28;582(12):1667-1671.
- ¹⁸¹ Sen CK, Sashwati R, Packer L. Fas mediated apoptosis of human Jurkat T-cells: intracellular events and potentiation by redox-active alpha-lipoic acid. *Cell Death Differ.* 1999 May;6(5):481-91.
- ¹⁸² Wenzel U, Nickel A, Daniel H. Alpha-lipoic acid induces apoptosis in human colon cancer cells by increasing mitochondrial respiration with a concomitant O₂-generation. *Apoptosis.* 2005 Mar;10(2):359-68.
- ¹⁸³ Simbula G, Columbano A, et al. Increased ROS generation and p53 activation in alpha-lipoic acid-induced apoptosis of hepatoma cells. *Apoptosis.* 2007 Jan;12(1):113-23.
- ¹⁸⁴ Berkson BM, Rubin DM, Berkson AJ. The long-term survival of a patient with pancreatic cancer with metastases to the liver after treatment with the intravenous alpha-lipoic acid/low-dose naltrexone protocol. *Integr Cancer Ther.* 2006 Mar;5(1):83-9.
- ¹⁸⁵ Berger V, et al. Effect of thioctic acid on the chemotherapeutic efficacy of cyclophosphamide and vincristine sulfate. *Arzneimittelforschung.* 1983;33(9):1286-1288.
- ¹⁸⁶ Casciari JJ, Riordan NH, Schmidt TL, et al. Cytotoxicity of ascorbate, lipoic acid, and other antioxidants in hollow fibre in vitro tumours. *Br J Cancer.* 2001 Jun 1;84(11):1544-1550.
- ¹⁸⁷ Van de Mark K, Chen JS, Steliou K, Perrine SP, Faller DV. Alpha-lipoic acid induces p27Kip-dependent cell cycle arrest in non-transformed cell lines and apoptosis in tumor cell lines. *J Cell Physiol.* 2003 Mar;194(3):325-40.
- ¹⁸⁸ Ho YS, Lai CS, Liu HI, et al. Dihydrolipoic acid inhibits skin tumor promotion through anti-inflammation and anti-oxidation. *Biochem Pharmacol.* 2007 Jun 1;73(11):1786-95.
- ¹⁸⁹ Vig-Varga E, Benson EA, Limbil TL, et al. Alpha-lipoic acid modulates ovarian surface epithelial cell growth. *Gynecol Oncol.* 2006 Oct;103(1):45-52.
- ¹⁹⁰ Karpov LM, Dvuzhil'naia ED, Savvov VI, Phan Van Thuy. [S35 lipoic acid distribution and its effect on pyruvate dehydrogenase activity in rats with Walker carcinoma] *Vopr Onkol.* 1977;23(10):87-90.

¹⁹¹ Wallace LA, Pellizzari ED, Hartwell TD, et al. The TEAM (Total Exposure Assessment Methodology) Study: personal exposures to toxic substances in air, drinking water, and breath of 400 residents of New Jersey, North Carolina, and North Dakota. *Environ Res.* 1987 Aug;43(2):290-307.